



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07K 13/00, C07H 15/12 A61K 37/02	A1	(11) International Publication Number: WO 93/24529 (43) International Publication Date: 9 December 1993 (09.12.93)
(21) International Application Number: PCT/US93/05358 (22) International Filing Date: 4 June 1993 (04.06.93) (30) Priority data: 07/894,215 4 June 1992 (04.06.92) US (71) Applicant: UNIVERSITY OF SOUTHERN CALIFORNIA [US/US]; University Park, Los Angeles, CA 90089 (US). (72) Inventors: JOHNSON, Lincoln, V. ; 411 Woodcliffe Road, Pasadena, CA 91105 (US). TOMBRAN-TINK, Joyce ; 8510 16th Street, #409, Silverspring, MD 20910 (US). (74) Agent: SHARP, Janice, A.; Christie, Parker & Hale, 350 W. Colorado Blvd., 5th Fl., P.O. Box 7068, Pasadena, CA 91109-7068 (US).		(81) Designated States: AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: RETINAL PIGMENTED EPITHELIUM DERIVED NEUROTROPHIC FACTOR (57) Abstract The present invention relates to a purified retinal pigmented epithelium derived neurotrophic factor composition and a method for purifying such a retinal pigmented epithelium neurotrophic factor. The present invention also relates to a recombinant DNA molecule comprising a gene encoding a retinal pigmented epithelium derived neurotrophic factor having the DNA sequence or the amino acid sequence in SEQ ID NO 1 and to an organism transformed with the recombinant DNA molecule. In addition, the present invention relates to a method of treating tumors, ocular diseases, nerve injuries, and conditions resulting from the activity of serine proteases, which comprises administering retinal epithelium derived neurotrophic factor.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

1

5

10

-1-

RETINAL PIGMENTED EPITHELIUM DERIVED NEUROTROPHIC FACTOR

15 This invention was made with government support under grant EY04741, awarded by
The National Institutes of Health. The United States government has certain rights in this
invention.

Field of the Invention

20 This invention relates to the purification and use of a retinal pigmented epithelium
derived neurotrophic factor (PEDNF).

Background of the Invention

Many types of neurons depend upon the availability of special regulatory molecules,
known as neurotrophic factors, for their survival and well-being. The best characterized of
25 the neurotrophic factors is nerve growth factor (NGF). NGF regulates the survival and
specialized function of sympathetic and dorsal root ganglion neurons in the peripheral nervous
system and of some cholinergic neurons in the central nervous system. Trophic factors,
which act on other neurons, have also been identified, and two such factors, ciliary
neurotrophic factor (CNTF) and brain-derived neurotrophic factor (BDNF) have been
30 purified. Moreover, it has recently been shown that some growth factors, such as fibroblast
growth factor (FGF) and epidermal growth factor (EGF), which initially were identified based
on their mitogenic effects upon cells, also function as survival-promoting agents for some
neurons. Post-synaptic target cells and satellite cells, such as glia cells, appear to be major
sources of neurotrophic factors.

35 It has been proposed that the survival of retinal photoreceptor cells may also be
regulated by specific neurotrophic factors. Evidence supporting this concept includes the
observation that photoreceptors undergo developmental neuronal death in some species, a

-2-

1 phenomenon which is generally considered to reflect the limited availability of neurotrophic factors. Photo-receptor development, as well as maintenance of normal function, has also been shown to require interactions with the retinal pigment epithelium (RPE), suggesting that RPE-derived molecules or factors could be necessary for photoreceptor function and survival.

5 The RPE develops in advance of and lies adjacent to the neural retina. A closed compartment between the two cell layers contains the interphotoreceptor matrix, and many soluble secretory products of RPE and neural retina cells are contained in the interphotoreceptor matrix. Nutrients, metabolites or trophic factors exchanged between the RPE and neural retina, must pass through the interphotoreceptor matrix. RPE cells, for
10 example, are thought to synthesize and secrete a photoreceptor survival-promoting factor (PSPA).

Cultured RPE cells synthesize a number of well known trophic factors, including platelet derived growth factor (PDGF), FGF, transforming growth factor- α (TGF- α), and transforming growth factor- β (TGF- β). It is possible that these or other unknown factors
15 derived from RPE could influence the development of the neural retina.

The neural-derived RPE forms a monolayer of cells interposed between the neural retina and circulating blood within the choroid. In this strategic location, the RPE forms a part of the blood-retina barrier, performs functions essential to retinal integrity and functions, and plays important roles in vascular, inflammatory, degenerative, and dystrophic diseases
20 of the retina and choroid. The functions of the RPE in relation to the visual process are several-fold and include light-dark adaption, phagocytosis of shed rod outer segment membrane and nutrition. On the other hand, the close interdependence of the RPE and the neural retina during normal development have been known for a long time, but functionally are not well understood, although it is known that the RPE is important for retinal
25 regeneration. It has been consistently observed that loss of contact of the neural retina with the RPE of many vertebrates (retinal detachment) results in degeneration of the retina. As a side effect of the retinal detachment, strong cell proliferation, originating from the RPE which underlies the spots of detachment, has often been observed.

Thus, identification of hypothetical survival-promoting factors for photoreceptor cells
30 would potentially be of great importance for the treatment of pathological conditions which result in blindness due to photoreceptor degeneration of unknown etiology. While these types of selective photoreceptor degenerations could be due to a variety of different mechanisms, analogies with neuronal degenerations in other regions of the nervous system suggest the possible involvement of a neurotrophic activity in the retina.

35

1 Summary of the Invention

5 The present invention relates to a purified retinal pigmented epithelium derived neurotrophic factor composition and a method for purifying such a retinal pigmented epithelium neurotrophic factor. The purification procedure comprises providing an impure protein fraction containing retinal pigmented epithelium derived neurotrophic factor and applying the impure protein fraction containing retinal pigmented epithelium derived neurotrophic factor to a cation-exchange chromatography medium. The cation-exchange chromatography medium is then washed to elute any unbound proteins and the retinal pigmented epithelium derived neurotrophic factor is eluted from the cation-exchange chromatography medium and collected.

10 The present invention also relates to a recombinant DNA molecule comprising a gene encoding a retinal pigmented epithelium derived neurotrophic factor having the DNA sequence or the amino acid sequence in SEQ ID NO 1 and to an organism transformed with a recombinant DNA molecule comprising a retinal pigmented epithelium derived neurotrophic factor gene having a DNA sequence identified in SEQ ID NO 1.

15 The present invention also relates to a method of treating tumors, ocular diseases and conditions resulting from the activity of serine proteases which comprises administering PEDNF.

20

25

30

35

1 Brief Description of the Drawings

 Features and advantages of the invention will be more apparent from a reading of the claims and of the detailed description of the invention in conjunction with the drawings described below.

5 Fig. 1 is a Coomassie blue stained SDS polyacrylamide gel showing the location of the PEDNF protein doublet unique to RPE-conditioned medium (RPE-CM).

 Fig. 2 is a differential interference contrast micrograph showing cell cultures 8-day post-attachment stimulated for 7 days in serum-free medium containing 2 μ g/ml of electroeluted PEDNF (panel A) or serum-free control medium (panel B).

10 Fig. 3 is a silver-stained, two-dimensional gel. Molecular weights are indicated at the left, and pH extremes are indicated at the top.

15

20

25

30

35

1 Detailed Description

Cells of the retinal pigmented epithelium are closely associated with differentiating retinoblasts in vivo and contribute to an environment essential to their development and normal function, both in vivo and in vitro. Retinoblastoma cells exhibit a multipotential
5 differentiative nature paralleling that of their precursor, the primitive retinoblast, as evidenced by the fact that agents, such as laminin, sodium butyrate and dibutyl cAMP, can induce the expression of neuronal, glial, and pigmented epithelial characteristics in vitro.

A human retinoblastoma cultured cell line, Y79 cells, when exposed to RPE-conditioned medium (RPE-CM), exhibits a high degree of neuronal-like differentiation. The
10 differentiation is observed in both the morphological and biochemical characteristics of the cells. The exposed cells extend arborizing neuritic processes from their cell bodies and express elevated levels of neuron-specific enolase (NSE) and neurofilament proteins.

RPE-secreted proteins have been fractionated from RPE-CM, and a protein doublet, with an apparent molecular weight of about 50,000 to about 55,000, that is unique to RPE-
15 CM has been identified. This pigment epithelium derived neurotrophic factor (PEDNF), which is a major secretory product of human fetal RPE cells, has been isolated and shown to have neurotrophic effects on Y79 retinoblastoma cells.

PEDNF has uses in the treatment of retinal diseases including, but not limited to, retinoblastoma and other ocular tumors, retinitis pigmentosa, various forms of retinal
20 detachment, macular degeneration, diabetic retinopathy, and other inherited and age-related pathologies of retinal cells.

In the case of retinal tumors, PEDNF induces the tumor cells to display biochemical and phenotypic characteristics of mature neuronal cells. Such changes are identified by a
25 cessation or reduction in the rate of cell division, which leads to tumor regression or a slowing in the rate of tumor growth. In the case of non-tumorous retinal diseases, the neurotrophic properties of PEDNF enhance survival and well-being of photoreceptor or other retinal cells, prolonging their functional life span and delaying the rate of the onset of impaired vision and ultimate blindness. In the case of retinal detachment, PEDNF prolongs the life span of photoreceptor cells sufficiently to allow standard reattachment procedures to
30 be effective in re-establishing the retina-RPE interface, thereby restoring normal vision.

1. Preparation of an Impure Protein Fraction Containing Retinal
Pigmented Epithelium Derived Neurotrophic Factor

Retinal pigmented epithelium derived neurotrophic factor (PEDNF) may be isolated
35 from any tissue or cell producing PEDNF.

Naturally-occurring cells that produce PEDNF are retinal pigmented epithelium cells. Such cells may be grown in culture to secrete PEDNF into the medium in which they are

-6-

1 growing. The medium may be harvested periodically, and the PEDNF isolated from the media. Suitable cell cultures may be established from retinal pigmented epithelium cells derived from humans, monkeys, and other primates or other animals, such as chickens, mice, rats, cows and pigs.

5 PEDNF may also be isolated by extraction of the interphotoreceptor matrix (IPM) or from the retina of humans, monkeys, and other primates or other animals, such as chickens, mice, rats, cows and pigs.

Alternatively PEDNF may be derived from sources in which it is not naturally-occurring, such as organisms transfected with a recombinant DNA molecule constructed to result in the expression of PEDNF in the host cells chosen for the expression of the gene.

A. Culturing of Human Retinal Pigmented Epithelium (RPE) Cells

15 Cultures of RPE cells are established by harvesting post-mortem eyes by aseptically opening the eyes and removing the vitreous body and retina. The exposed RPE is washed with a buffered solution such as modified Earle's balanced salt solution (MEBS: 115.5 mM NaCl, 3.5 mM KCl, 1 mM NaH_2PO_4 , 0.5 mM CaCl_2 , 0.27 mM MgCl_2 , 0.37 mM MgSO_4 , 15 mM HEPES, 14 mM NaHCO_3 , 12 mM glucose, pH 7.2). RPE cells are scraped from Bruch's membrane or are dislodged by a stream of fluid such as MEBS or cell culture medium, applied using a Pasteur or similar pipette. This step may be preceded by exposure to proteolytic or other enzyme(s), such as Dispase (Boehringer-Mannheim, Indianapolis, IN, Catalog #295 825). Alternatively, RPE cells may be isolated by detaching sections of sclera from an intact eye to expose choroidal tissue and treating this choroidal surface with proteolytic or other enzymes such as Dispase prior to removal of the vitreous and retina and dislodging RPE cells by the spraying of cell culture medium as described by Pfeffer. Tissue fragments are transferred to tissue culture dishes in a medium such as "low Ca^{++} " or "high Ca^{++} " complete RPE-47 medium, as described by Pfeffer, or Eagles's minimal essential medium (MEM, supplied by GIBCO of Grand Island, NY) supplemented with about 0.5% to about 20%, by volume, fetal calf serum. At concentrations below about 0.5% fetal calf serum, the serum concentrations are too low to effectively support the growth of the cells. At concentrations above about 20% serum, no additional benefit, with respect to cell growth, is conferred on the cells.

25 The medium may also be supplemented with anti-biotics and/or fungicides to prevent the growth of bacteria and/or fungi in the cultures. Antibiotics and fungicides suitable for use in the present invention are about 1,000 units/ml penicillin, about 100 $\mu\text{g/ml}$ streptomycin, about 0.25 μg amphotericin, and about 50 $\mu\text{g/ml}$ gentamicin, or other suitable

-7-

1 such agents known in the art. These antibiotics may be used individually or in combination, as desired.

The cells are incubated at about 37°C in an atmosphere of about 5% carbon dioxide. Retinal pigmented epithelium (RPE) cells attach to the surface of the dishes, proliferate, and
5 eventually form confluent monolayers.

When the cells have grown to confluence, about 3-7 days from the initial culturing of the cells, they are harvested, for example, by trypsinizing the cell monolayer, or by other methods known to those skilled in the art, and resuspending the cells in a medium such as MEM, supplied by GIBCO, supplemented with about 5% to about 20% fetal calf serum. A
10 portion of the cells are reseeded into sterile tissue culture flasks, after which time the cells are again grown in an atmosphere of about 5% CO₂ at about 37°C.

Alternatively, RPE cells may be isolated by removing the cornea, the 2 mm scleral ring, and the vitreous and neural retina from the eyes. The eyes are washed with calcium and magnesium-free Hanks Balanced Salt Solution (HBSS: 1.3 mM CaCl₂, 5 mM KCl, 0.3 mM
15 KH₂PO₄, 0.5 mM MgCl₂, 0.4 mM MgSO₄, 138 mM NaCl, 4 mM NaHCO₃, 0.3 mM Na₂HPO₄, 5.6 mM D-glucose and 0.03 mM Phenol Red), supplied by GIBCO/BRL of Gaithersburg, MD. The eye cup is filled with a solution comprising about 0.1%, by weight, trypsin, about 0.1%, by weight, hyaluronidase in calcium and magnesium-free Hanks balanced salt solution, and the eye is incubated at about 37°C for about 15 to about 30 min.

20 The loose RPE cells are collected by gentle aspiration, and the procedure is repeated until the RPE cells are released. The trypsin is inactivated by adding about 5% to about 20% fetal calf serum to the cell sample. The cells are collected by centrifugation at about 1,200 rpm for about 7 min.

The RPE cells are then plated onto sterile tissue culture plates at a density of about
25 1 x 10⁵ cells for a 35 mm plate. (Proportionally more or less cells are plated if larger or smaller plates or containers are used.) The cells are grown in DulBecco's Modified Eagles Medium (DMEM), supplied by GIBCO, or other suitable medium. The medium may be supplemented with an equal volume of HAM's F12 medium (supplied by GIBCO), about 1 mM sodium pyruvate, about 0.625 mM Hepes, about 6 mM L-glutamine, about 1% non-
30 essential amino acids, about 5 µg/ml insulin, about 5 µg/ml transferrin, about 5 ng/ml selenium, antibiotics as described above, and about 0.5% to about 20% fetal calf serum, as described above. The insulin, transferrin, and selenium are supplied by Collaborative Research of Lexington, MA. Other reagents suitable for use in the present invention, unless otherwise specified, are supplied by Sigma Chemical Co. of St Louis, MO.

35 When the cells have grown to confluence (about 3-7 days from the initial culturing of the cells), they are harvested, for example, by trypsinizing the cell monolayer or by other methods known to those skilled in the art, and resuspending the cells in a medium such as

-8-

1 MEM supplemented with about 0.5% to about 20% fetal calf serum. A portion of the cells are reseeded into sterile tissue culture flasks, after which time the cells are again grown in an atmosphere of about 5% CO₂ at about 37°C.

5 While only two methods for the culturing of RPE cells are described, other suitable methods for culturing RPE cells are known in the art. Such methods are also suitable for use in the practice of the present invention.

B. Preparation of Retinal Pigmented Epithelium
Conditioned Medium (RPE-CM)

10 PEDNF is a secreted protein, and RPE cells secrete PEDNF into the medium in which they are grown. Therefore, a convenient method of producing PEDNF is to grow RPE cells in culture and to periodically harvest the media from the cultures.

15 A method suitable for use in the present invention for isolating PEDNF from medium is to allow RPE cells to grow to confluence in a medium such as DMEM supplemented with about 1 mM sodium pyruvate, about 0.625 mM Hepes, about 6 mM L-glutamine, about 1% non-essential amino acids, about 5 µg/ml insulin, about 5 µg/ml transferrin, about 5 ng/ml selenium, antibiotics and/or fungicides as described above, and about 0.5% to about 20% fetal calf serum, as described above. The confluent cultures of RPE cells are washed extensively with Hank's balanced salt solution, or other suitable wash solutions, to remove serum proteins derived from the fetal calf serum present in the media which is used to culture the RPE cells. About 1 ml, per cm² of cell surface area, of serum-free medium such as DMEM supplemented with about 1 mM sodium pyruvate, about 0.625 mM Hepes, about 6 mM L-glutamine about 1% non-essential amino acids, about 5 µg/ml insulin, about 5 µg/ml transferrin, about 5 ng/ml selenium, and antibiotics and/or fungicides as described above, is added to the cultures, and they are incubated in an atmosphere of about 5% CO₂ at about 37°C for about 1 to about 10 days. Alternatively, PEDNF can be isolated from serum-containing medium such as DMEM supplemented with about 1 mM sodium pyruvate, about 0.625 mM Hepes, about 6 mM l-glutamine, about 1% non-essential amino acids, about 5 µg/ml insulin, about 5 µg/ml transferrin, about 5 ng/ml selenium, antibiotics and/or fungicides as described above, and about 0.5% to about 20% fetal calf serum as described above.

25 The medium is then collected by pouring the medium into plastic centrifuge tubes, and the medium is centrifuged at about 3,000 rpm for about 10 min. to remove any free cells and other particulate matter from the medium, and filtered to provide an impure PEDNF protein solution. The medium may be used directly for the purification or testing of PEDNF or it may be stored at -20°C until required.

35

1 C. Isolation of PEDNF from Tissue Samples

 PEDNF may be isolated directly from eyes by opening the eyes and removing the vitreous body and retina. The retinal pigmented epithelium is scraped off the Bruch's membrane using a disposable cell scraper. Tissue fragments are transferred to a Teflon or glass homogenizer in 10 mM phosphate-buffered saline (PBS) and homogenized to break the cells. The solution is then centrifuged at about 10,000 rpm for about 10 min. and filtered to remove cell debris. The supernatant, which contains PEDNF, is collected to provide an impure PEDNF protein solution. The medium may be used directly for the purification or testing of PEDNF or it may be stored at -20°C until required.

10

 D. Isolation of PEDNF from Recombinant Cells

 PEDNF may also be isolated from recombinant cells which have been constructed to express the PEDNF gene. The PEDNF may be expressed as an intracellular or an extracellular protein.

15

 Intracellular PEDNF is isolated by homogenizing the cells in a Dounce or other suitable homogenizer in a buffer, such as PBS, that may contain detergents or other solubilizing agents such as urea or guanidine hydrochlorid, and centrifuging at about 10,000 rpm for about 20 min. to remove cellular debris, to provide an impure PEDNF protein fraction.

20

 Extracellular PEDNF is isolated by collecting the medium in which cells expressing PEDNF are grown. This is most conveniently performed in a continuous centrifugation process, such as with a Sharples centrifuge. The supernatant is collected to provide an impure PEDNF protein fraction. The medium may be used directly for the purification or testing of PEDNF or it may be stored at -20°C until required.

25

2. Purification of PEDNF

 A. Small Scale Purification of PEDNF

 i. Ammonium Sulfate Precipitation

 An impure PEDNF protein fraction may be partially purified by ammonium sulfate precipitation to provide an ammonium sulfate purified PEDNF protein fraction. Percent ammonium sulfate refers to % saturation of ammonium sulfate at 20°C and is based on a 100% saturation of 767 g/l.

 An impure PEDNF protein fraction is brought to about 50% saturation with ammonium sulfate by the addition of about 313 g of solid ammonium sulfate per liter of impure protein fraction at about 20°C. The ammonium sulfate is preferably added slowly to the impure protein fraction while the solution is stirred, such as with a stir bar on a mechanical stirrer. The ammonium sulfate is added slowly to prevent localized high

35

-10-

1 concentrations of ammonium sulfate that may result in rapid precipitation and denaturation
of proteins present in the impure protein fraction. After all the ammonium sulfate is added,
the about-50% ammonium sulfate solution is stirred for about 30 min. at about 20°C. The
about-50% ammonium sulfate solution is then centrifuged at about 10,000 g, at about 20°C
5 for about 20 min. The supernatant is collected for further processing. Alternatively, the
about-50% ammonium sulfate solution may be filtered through filter paper such as Whatman
#1, to remove the precipitate. In this case, the filtrate is collected for further processing.

The about-50% ammonium sulfate solution is then brought to about 70% saturation
by the addition of about 137 g/l of ammonium sulfate. The ammonium sulfate is added
10 slowly, and after all the ammonium sulfate is added, the about-70% ammonium sulfate
solution is stirred for about 30 min. at about 20°C. The about-70% ammonium sulfate
solution is centrifuged or filtered, as described above, and the precipitate is collected.

The ammonium sulfate precipitate is then redissolved in a buffer such as about 10 mM
phosphate, pH 7, to form a 50-70% ammonium sulfate fraction. The solution is then
15 diafiltered using an Amicon Diaflo ultrafiltration unit, or dialyzed against a buffer such as
about 10 mM phosphate, pH 7, to remove the residual ammonium sulfate from the
redissolved 50%-70% ammonium sulfate fraction.

20 ii. Purification of PEDNF by SDS Gel
Electrophoresis

A small-scale isolation of PEDNF is conducted by SDS-polyacrylamide slab gels.
Such polyacrylamide gels are well known in the art, and the preparation of such gels has been
described by Weber and Osborn, J. Biol. Chem., 244, 4406 (1969), as modified by Laemmli,
Nature, 227, 680 (1970).

25 Samples of an impure PEDNF protein fraction or ammonium sulfate purified PEDNF
protein fraction are dialyzed against a buffer such as about 10 mM sodium phosphate buffer,
pH 7.5, lyophilized and resuspended in 62.5 mM Tris, pH 6.8, 2% SDS, 10% glycerol,
0.001% bromophenol blue, and 0.1 M 2-mercaptoethanol, wherein the bromophenol blue is
a migration marker. Additional samples to be loaded on the gel include molecular-weight
standards such as phosphorylase B, bovine serum albumin, ovalbumin, carbonic anhydrase,
30 soybean trypsin inhibitor and lysozyme (such as supplied by Bio-Rad, catalog #161-0304),
and controls as may be necessary. In cases where conditioned media are used as samples,
media which have not been exposed to RPE cells (unconditioned media) are included as
controls.

35 The samples are then incubated in a boiling-water bath for about 5 min. and loaded
onto SDS-polyacrylamide slab gels. The markers and unconditioned media are loaded into
wells on the outside of the gel, and the impure PEDNF protein fractions or the ammonium

-11-

1 sulfate purified PEDNF protein fraction are loaded on the remaining inside wells. The
samples are then subjected to electrophoresis. The electrophoresis is continued until the
bromphenol blue marker has reached the bottom of the gel. At the completion of
electrophoresis, strips from each outside edge of the gels, which included the markers and
5 a lane each of an impure PEDNF protein fraction and an unconditioned media control, are
stained with Coomassie blue. The stained protein bands which develop on the stained strips
are aligned with the unstained portion of the gel to locate the position of the proteins present
in the impure PEDNF protein fraction or the ammonium sulfate purified PEDNF protein
fraction but absent from control media.

10 A PEDNF protein doublet, with an apparent molecular weight of about 50,000 to
about 55,000, unique to the impure PEDNF protein fraction, is excised and the proteins
electroeluted, by methods known in the art, from the unstained portion of the gel. The eluant
is centrifuged to remove gel fragments and the supernatant dialyzed against 10 mM
phosphate-buffered saline (145 mM NaCl, 8.1 mM Na_2HPO_4 , and 1.9 mM $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$)
15 to provide an SDS-polyacrylamide purified PEDNF protein fraction.

iii. Purification of PEDNF by
Cation-Exchange HPLC

20 An impure PEDNF protein fraction or an ammonium sulfate purified PEDNF protein
fraction is dialyzed against water or a buffer such as about 10 mM phosphate buffer, pH 7.2,
or other suitable buffer, to remove media and salts from the impure protein sample. PEDNF
may be purified by cation-exchange HPLC using a column chromatography medium, such as
that supplied under the trade name "Brownlee Aquapore CX-300" by Western Analytical,
Temecula, CA, packed into a column, such as a 4.6 x 30 mm column, or other suitable
25 cation-exchange HPLC chromatography medium.

The chromatography medium is equilibrated with a buffer such as about 10 mM
phosphate, pH 7.2. The dialyzed impure PEDNF protein fraction or ammonium sulfate
purified PEDNF protein fraction is loaded onto the chromatography medium, and the
chromatography medium with PEDNF bound to it is washed with a buffer such as about 10
30 mM phosphate, pH 7.2, until all unbound proteins are washed from the chromatography
medium. PEDNF is eluted from the chromatograph medium with a linear salt gradient from
about 0.0 to about 0.5 M NaCl. PEDNF elutes as a single peak with a NaCl concentration
of about 0.25 M.

35 The eluted PEDNF is concentrated by lyophilization and resolubilized in water or a
buffer such as about 10 mM phosphate buffer, pH 7.2, or other suitable buffer, to form a
cation-exchange HPLC purified PEDNF protein fraction.

1 iv. Purification of PEDNF by
 Reverse-Phase HPLC

5 An impure PEDNF protein fraction or ammonium sulfate purified PEDNF protein fraction is dialyzed against a buffer such as about 10 mM phosphate buffer, pH 7.2, or other suitable buffer, to remove media and salts from the impure protein sample. PEDNF may be purified by reverse-phase HPLC using a column chromatography medium such as a chromatography medium supplied under the trade name "Vydac C8" by The Separation Group of Hesperia, CA, packed into a column such as a 4.6 x 250 mm column or other suitable reverse-phase HPLC chromatography medium.

10 The chromatography medium is equilibrated with a solution such as about 0.1%, by volume, trifluoroacetic acid (TFA). The dialyzed impure PEDNF protein fraction or ammonium sulfate purified PEDNF protein fraction is loaded onto the chromatography medium, and the chromatography medium with PEDNF bound to it is washed with an eluant such as 0.1%, by volume, TFA, until all unbound proteins are washed from the chromatography medium. PEDNF is eluted from the chromatograph medium with a linear gradient from about 0.1% TFA in water to about 95% acetonitrile (CH₃CN), 0.1% TFA, 5% H₂O. PEDNF elutes as a single peak with a CH₃CN concentration of about 70%.

15 The eluted PEDNF is concentrated by lyophilization and resolubilized in water or a buffer such as about 10 mM phosphate buffer, pH 7.2, or other suitable buffer, to form a reverse-phase HPLC purified PEDNF protein fraction.

20 v. Size-Exclusion HPLC

25 An impure PEDNF protein fraction, an ammonium sulfate purified PEDNF protein fraction, a cation-exchange HPLC purified PEDNF protein fraction, or a reverse-phase HPLC purified PEDNF protein fraction may be purified by size-exclusion chromatograph using a chromatography medium, such as that supplied under the trade name "Bio-Rad TSK-250" by Bio-Rad of Richmond, CA, packed into a column such as a 7.5 X 300 mm column. The impure PEDNF protein fraction, the ammonium sulfate purified PEDNF protein fraction, the cation-exchange HPLC purified PEDNF protein fraction, or the reverse-phase HPLC purified PEDNF protein fraction is loaded onto the size-exclusion chromatography medium, which has been equilibrated with a buffer such as 0.02 M Tris-HCl, pH 7.0, 0.6 M NaCl. PEDNF-containing fractions are collected and dialyzed against a buffer such as about 10 mM phosphate buffer, pH 7.2 to provide a size-exclusion purified PEDNF protein fraction.

35

1 vi. Heparin Chromatography

 An impure PEDNF protein fraction or an ammonium sulfate purified PEDNF protein fraction may also be purified by heparin chromatography. A heparin chromatography medium such as heparin agarose, supplied by Sigma Chemical Co. of St Louis, MO (Cat. 5 No. H-5380), is equilibrated with a buffer such as about 10 mM Tris-HCl, pH 7.5.

 An impure PEDNF protein fraction or an ammonium sulfate purified PEDNF protein fraction is dialyzed against a buffer such as about 10 mM Tris, pH 7.5, to remove any salts or media from the samples, and the dialyzed PEDNF solution is applied to the equilibrated heparin agarose. After the PEDNF solution has been applied to the heparin agarose, the 10 heparin agarose is washed with a buffer such as about 10 mM Tris-HCl, pH 7.5, until all unbound proteins are eluted from the heparin agarose. PEDNF is then eluted from the heparin agarose with a buffer such as about 10 mM Tris-HCl, pH 7.5, 0.5 M NaCl.

 The eluate containing PEDNF is then diafiltered in an Amicon Diaflo ultrafiltration unit, or is dialyzed against a buffer such as about 10 mM Tris-HCl, pH 7.5, to remove NaCl 15 present in the eluate, thereby providing a heparin purified PEDNF protein fraction.

 The above-described purification procedures may use an impure PEDNF protein fraction or an ammonium sulfate purified PEDNF protein fraction as the starting material for the subsequent column purification. Alternatively, a protein fraction which has already been purified by one or more of the described chromatography steps could also be used. 20 Therefore, the purification procedures may be used alone or in combination with each other or with other purification techniques known in the art to produce a PEDNF protein fraction of the desired purity.

 B. Large-Scale Preparation of PEDNF

25 i. Ammonium Sulfate Precipitation

 Large-scale purification of PEDNF may be by ammonium sulfate precipitation to provide an ammonium sulfate purified PEDNF protein fraction.

 An impure PEDNF protein fraction is brought to about 50% saturation with ammonium sulfate by the addition of about 313 g of solid ammonium sulfate per liter of 30 impure PEDNF protein fraction. The ammonium sulfate is preferably added slowly to the impure protein fraction while the solution is stirred, such as with a stir bar on a mechanical stirrer. The ammonium sulfate is added slowly to prevent localized high concentrations of ammonium sulfate which may result in rapid precipitation, and denaturation of proteins present in the impure protein fraction. After all the ammonium sulfate is added, the 50% ammonium sulfate solution is stirred for about 30 min. at 20°C. The 50% ammonium sulfate 35 solution is then centrifuged at about 10,000 g, at 20°C for about 20 min. The supernatant is collected for further processing. Alternatively, the 50% ammonium sulfate solution may

1 be filtered through filter paper, such as Whatman #1, to remove the precipitate. The filtrate is collected for further processing.

5 The 50% ammonium sulfate solution is then brought to about 70% saturation by the addition of about 137 g/l of ammonium sulfate. The ammonium sulfate is added slowly, and after all the ammonium sulfate is added, the 70% ammonium sulfate solution is stirred for about 30 min. at 20°C. The 70% ammonium sulfate solution is centrifuged or filtered, as described above, and the precipitate is collected.

10 The ammonium sulfate precipitate is then redissolved in a buffer such as about 10 mM phosphate, pH 7, to form a 55-70% ammonium sulphate fraction, then diafiltered using an Amicon Diaflo ultrafiltration unit, or dialyzed against about 10 mM sodium phosphate, pH 7, to remove the ammonium sulfate from the redissolved 55-70% fraction to provide an ammonium sulfate purified PEDNF protein fraction.

15 ii. Anion-Exchange Chromatography

The pI of the PEDNF protein is about 3.9 to about 7.2. Therefore, at a pH of about 7.5, the protein has a net negative charge and binds to an anion-exchange chromatography medium such as DEAE (diethylaminoethyl) cellulose.

20 For use, an anion-exchange chromatography medium such as DEAE cellulose is equilibrated with a buffer such as about 10 mM Tris, pH 7.5. The PEDNF is applied to the anion-exchange chromatography medium, and the medium is washed with a buffer such as about 10 mM Tris, pH 7.5, until all unbound proteins are eluted from the column. After all the unbound proteins are eluted, PEDNF is eluted with a linear salt gradient from about 0 to about 1 M NaCl in a buffer such as about 10 mM Tris-HCl, pH 7.5. The fractions containing PEDNF are collected and pooled. These fractions are then diafiltered or dialyzed
25 against a buffer such as about 10 mM Tris-HCl, pH 7.5, to remove NaCl, thereby providing an anion-exchange chromatography purified PEDNF protein fraction.

Anion-exchange chromatography may be conducted by either batch or column chromatography techniques.

30 iii. Heparin Chromatography

Heparin chromatography may also be used for the large-scale purification of PEDNF. A heparin chromatography medium such as heparin agarose, supplied by Sigma Chemical Co. (Cat. No. H-5380), is equilibrated with a buffer such as about 10 mM Tris-HCl, pH 7.5.

35 An impure PEDNF protein fraction or an ammonium sulfate purified PEDNF protein fraction is dialyzed against a buffer such as about 10 mM Tris-HCl, pH 7.5, and then applied to the heparin agarose. After the PEDNF solution has been applied to the heparin agarose, the heparin agarose is washed with a buffer such as about 10 mM Tris-HCl, pH 7.5, until all

-15-

1 unbound proteins are eluted from the heparin agarose. PEDNF is then eluted from the heparin agarose with a buffer such as about 10 mM Tris-HCl, pH 7.5, 0.5 M NaCl.

The eluant containing PEDNF is collected and diafiltered in an Amicon Diaflo ultrafiltration unit, or dialyzed against a buffer such as about 10 mM Tris-HCl, pH 7.5, to
5 remove the NaCl present in the eluate, to provide a heparin purified PEDNF protein fraction.

The above-described purification procedures may use an impure PEDNF protein fraction or an ammonium sulfate purified PEDNF protein fraction as the starting material for the subsequent column purification. Alternatively, a protein fraction which has already been purified by one or more of the described chromatography steps could also be used.
10 Therefore, the purification procedures may be used alone or in combination with each other or with other purification techniques known in the art to produce a PEDNF protein fraction of the desired purity.

3. PEDNF Assays

15 A. SDS Gel Electrophoresis

PEDNF may be identified by SDS-polyacrylamide gel electrophoresis on, for example, 7.5%, 10%, 12.5% SDS-polyacrylamide gels. The preparation of such gels has been described by Weber and Osborn, as modified by Laemmli, and the preparation and use of such gels are well known in the art.

20 The PEDNF protein samples are mixed with about 5 μ l of 62.5 mM Tris, pH 6.8, 12% SDS, 0.001% bromophenol blue, 10% glycerol, and 0.1 M 2-mercaptoethanol, wherein the bromophenol blue is a marker. Molecular-weight marker samples which include molecular-weight standards such as phosphorylase B, bovine serum albumin, ovalbumin, carbonic anhydrase, soybean trypsin inhibitor and lysozyme (such as supplied by Bio-Rad,
25 catalog #161-0304) are included as controls. The PEDNF protein samples and the molecular-weight standards are boiled for about 5 min. and loaded onto SDS-polyacrylamide slab gels. The samples are then subjected to electrophoresis until the bromophenol blue has migrated to the bottom of the gel. At the completion of electrophoresis, the gel is silver-stained, stained with Coomassie blue, or stained by other suitable protein staining methods. The molecular
30 weight of proteins in the PEDNF sample is then compared to the molecular-weight standards.

PEDNF migrates as a protein doublet, with an apparent molecular weight of from about 50,000 to about 55,000 on SDS polyacrylamide gels.

B. Neuronal Inductivity

35 PEDNF activity in protein samples may be assayed by its neuronal inductivity. Various concentrations of PEDNF are added to cultures of Y79 retinoblastoma (RB) cells, supplied by the American Type Culture Collection, Access No. HTB 18, of Rockville, MD.

-16-

1 The Y79 RB cells are grown in suspension culture. The cells are harvested by
centrifugation for about 5 min. at about 900 rpm at room temperature and resuspended in a
serum-free medium such as Dulbecco's modified Eagle's medium supplemented with 5 ug/ml
insulin, 5 ug/ml transferrin, 5 ng/ml selenous acid, and 876.6 ug/ml L-glutamine (serum-free
5 medium), which has previously been warmed to about 37°C. The collected cells are
resuspended at a concentration of about 10⁶ cells/ml in serum-free medium. About 50 to
about 500 ng/ml of PEDNF is added to about 25 ml aliquots of the cells. The cells are
incubated for about 7 days at about 37°C, then attached to poly-D-lysine-coated flasks or
glass coverslips. The poly-D-lysine-coated flasks are prepared by coating with a solution
10 containing about 200 ug/ml poly-D-lysine (such as that supplied by Sigma, Catalog #P7405)
for about 1-24 hours, followed by rinsing with water and serum-free medium.

i. Cell Analyses

15 Morphology of attached cells is monitored daily by phase contrast microscopy of living
cells using a microscope such as an inverted Diaphot TMD microscope, supplied by Nikon
of Tokyo, Japan, or by differential interference contrast microscopy of cells, fixed with a
fixative such as about 4% paraformaldehyde in 0.1 M sodium cacodylate buffer, using a
microscope, such as a BHS-BH2 microscope, supplied by Olympus of Tokyo, Japan.

20 Differentiation is assessed by calculating the percentage of cellular aggregates (more
than 90 % of Y79 cells plated from suspension culture attach to poly-D-lysine-coated flasks
as aggregates containing more than 5 cells) in which cells extend processes at day 1, 3, 7 and
11 after attachment. Experiments are performed in replicates of 3 and are repeated twice.

25 Expression of neuron-specific enolase (NSE) and neurofilament 200,000 molecular
weight protein subunit (NF 200) is monitored by immunofluorescence and viewed by either
epifluorescence microscopy (Olympus BHS-BH2 microscope) or microspectrofluorometry
(MSA, Farrand Microscope Spectrum Analyzer). Quantification is by 1) visual scoring of
intensity of fluorescence at 485 nm excitation as + = weak; ++ = moderate; +++ =
strong; ++++ = very intense, and 2) microspectrofluorometric analysis (MSA) readings
(uA x 100, time constant of about 0.3 seconds; specimen size of approximately 2 mm or
30 about 100 cells/aggregate; target size of about 15 µm or approximately that of 1 cell).

35 The presence of PEDNF in the protein sample results in the Y79 RB cells, extending
neurite-like processes. At a concentration of about 500 ng/ml, about 70% of the cells extend
neurite-like processes.

-17-

1 ii. Differentiation

Cells were cultured as described above. The cells were then monitored daily by phase-contrast microscopy of living cells (Olympus IMT-2) and by differential interference contrast microscopy (Olympus BHS) of cells fixed with a fixative such as about 4% paraformaldehyde. The percentage of differentiating cells is estimated by calculating the number of cellular aggregates containing five or more cells exhibiting neurite outgrowths following 8-10 days of culture on a poly-D-lysine substratum.

With about 50 to about 500 ng/ml PEDNF, approximately 80% of the cells undergo morphological differentiation within about 3 days.

10

4. Characterization of the PEDNF proteinA. Isolation of PEDNF Peptides

Purified PEDNF, about 500 μ g, is concentrated using Centricon 10 microconcentrators (Amicon, Danvers, MA), and then diluted in a suitable digestion buffer such as about 25 mM Tris, pH 8.5, 1 mM EDTA. To the protein sample is added a proteolytic enzyme, such as endoproteinase Lys-C, supplied by Boehringer-Mannheim of Indianapolis, IN. The PEDNF/proteinase mixture is incubated for about 18 hrs. at about 30°C, or until the reaction has gone to completion, i.e., until all the PEDNF is completely digested by the proteinase. Alternatively, the protein may be digested with trypsin in a buffer comprising about 10 mM PBS or by using other proteinases known in the art.

20

The resulting PEDNF polypeptide fragments are separated by using a separation system such as HPLC on a Vydac C8 reverse-phase column. A 4.6 X 250 mm column is suitable for use in the present invention. The column is equilibrated with about 0.1% TFA in water. The polypeptides are eluted with about 90% CH_3CN , 0.1% TFA and 5% H_2O .

25

Polypeptides eluted from the column which are well separated from other polypeptides are collected and subjected to protein sequencing analysis.

B. Protein Sequencing Analysis

The purified polypeptide fragments are subjected to amino-acid sequence analysis by methods known in the art. Alternatively, the amino-acid sequence analysis is conveniently performed under contract at an amino-acid sequencing facility, such as the Microsequencing Facility of Beckman Research Institute at the City of Hope in Duarte, CA.

30

35

1 5. Characterization of the PEDNF Gene

 A. Cloning of the PEDNF Gene

 Oligonucleotides are constructed from the sequence derived for the isolated polypeptide of PEDNF on an ABI 392 DNA/RNA Synthesizer or by methods well known in the art.

5 The oligonucleotides are used as primers for a polymerase chain reaction (PCR) by using a Techne thermal cycler and standard reagents and methodologies, supplied under the trade name "GeneAMP" by Perkin Elmer Cetus of Emeryville CA.

 A human fetal eye Charon BS cDNA library is screened by PCR techniques using a Techne thermal cycler and standard reagents and methodologies. The cDNA fragment generated by the reaction is isolated on a 3% NuSieve 3:1 gel (FMC Biochemicals) using NA-45 DEAE-cellulose paper (Schleicher and Schull) as described by Sambrook et al. 1989 In: "Molecular Cloning: A Laboratory Manual". 2nd ed. Cold Spring Harbor Press, Cold Spring Harbor, NY, which is incorporated herein by this reference. Briefly, the screening procedure is performed by taking about 1, 5 and 50 μ l aliquots of the library and placing them in 600- μ l siliconized reaction tubes and then bringing them to a final volume of about 74 μ l with double-distilled sterile water. The phage particles are disrupted by incubation at about 70°C for about 5 minutes and then cooling on wet ice.

 PCR master mix is made up in a 600- μ l reaction tube for 3 reaction tubes as follows:

 30 μ l of 10X Taq polymerase buffer;

20 24 μ l of dNTP mix; and an appropriate volume of double-distilled water is added to bring the master mix to a final volume of about 78 μ l.

 The final solution, therefore, comprises about 192 mM KCl, 38.5 mM Tris-HCl, pH 8.3, 51.8 mM MgCl₂, 0.038% (w/v) gelatin, 0.77 mM of each dNTP, and 3.8 μ l of each oligonucleotide primer. About 26 μ l of master mix is added to each reaction tube. 25 The library aliquots and the master mix solutions are overlaid with about 100 μ l of mineral oil and heated to about 94°C for about 5 min. The solutions are then equilibrated to the desired primer annealing temperature.

 After the solutions have been equilibrated to the desired primer annealing temperature, about 1.5 μ l of Taq polymerase is added to the solution and incubated for about 20 min.

30 Thermal cycling is continued by incubating: at about 72°C for about 3 min. for primer extension, however, this primer extension time may be varied if desired; at about 94°C for about 1 min. about 20 sec. to denature the extension product from the template; and at about 37°C for about 2 min. to anneal the primers to a template; and at about 72°C for about 3 min. for primer extension. The "cycle" is the repeated for a total of about 25 to about 30 35 cycles. At the end of the last cycle, a final primer extension step, of about 7 min. at 72°C is added.

-19-

1 The fragment is labelled with ^{32}P by random priming using a kit supplied under the trade name "Prime-It Random Primer Labeling Kit" by Stratagene. The labelled probe is used to screen about 200,000 plaque-forming units of the Charon BS cDNA library by methods well known in the art.

5 Positive clones are isolated, and the DNA purified, with reagents supplied under the trade name "Qiagen Maxi" by Qiagen, Inc. of Studio City, CA.

 The inserts within the phage vector are excised with an appropriate restriction enzyme, circularized with T4 DNA ligase supplied by New England Biolabs of Beverly, MA and transformed into competent *E. coli* Sure cells supplied by Stratagene, Inc. of La Jolla, CA.
10 The cells are then plated on Ampicillin/X-gal plates.

 White colonies are selected and mini-prepped using a Qiagen plasmid miniprep reagent supplied by Qiagen. Purified plasmids are digested to excise the insert, and the fragments are separated by gel electrophoresis to determine the size of the inserts. A clone with an appropriately-sized insert is then chosen for further investigation.

15 B. DNA Sequence Analysis

 Sequence analysis is performed with an automated sequencer or by other techniques well known in the art.

20 6. Expression of the PEDNF Gene in Recombinant Cells

 Commercial or large-scale production of PEDNF may be achieved by expression of the gene, after cloning into an appropriate vector, in a suitable host cell. One such suitable vector/host system is the baculovirus/insect cell systems.

 In one embodiment of the present invention, the PEDNF gene is cloned into a
25 baculovirus transfer vector such as pAc373, described by Lithgow et al., DNA and Cell Biology, 10, 443-449 (1991); pVL941, described by Ghiasi et al., Virology, 185, 187-194 (1991); or other such baculovirus transfer vectors that are well known by those skilled in the art.

 Generally, such vectors comprise the polyhedrin promoter of the Autographa californica nuclear polyhedrosis virus (AcNPV), inserted into a transfer vector such as pAc373, described by Summers and Smith (A manual for baculovirus vector and insect cell culture procedures, Texas Agric. Exp. Station Bull. No. 1555, which is incorporated herein by this reference). Recombinant plasmids are prepared by methods well known in the art, such as those described by Maniatis et al., "Molecular Cloning," Cold Spring Harbor
30 Laboratory, Cold Spring Harbor, NY (1982), which is incorporated herein by this reference.

35 The host cells, such as *S. frugiperda* (Sf9), are grown in TC100 medium supplied by GIBCO, supplemented with 10% fetal calf serum. The Sf9 cells are co-transfected with

-20-

1 purified infectious AcNPV DNA, about 1 μ g, and about 50 μ g of the recombinant DNA. Culture supernatants are harvested about 4 days after transfection. Recombinant viruses are identified by plaque hybridization or radiolabelling of the proteins produced.

5 The above description provides an example of the expression of the PEDNF gene in a vector/host expression system. Other expression systems are known in the art; for example, Saccharomyces cerevisiae has been used in conjunction with a number of vectors such as those described by Silar et al., Gene, 104, 99-102 (1991), Dietrich et al., Eur. J. Biochem., 201, 399-407 (1991), or Akiyoshi-Shibata et al., DNA and Cell Biology, 10, 613-612 (1991); recombinant vaccinia virus vectors in HeLa host cells such as those described by Nakano et al. Gene, 105, 173-178 (1991). Any such methods or other methods known in the art are suitable for use in the present invention for expression of the PEDNF gene.

10 The PEDNF gene may be expressed as a transported protein where the PEDNF is isolated from the medium in which the recombinant expression host is grown, or may be expressed as an intracellular protein by deleting the leader peptide, in which case the PEDNF is isolated from the host cells. The PEDNF so isolated is then purified by the methods described above.

7. Treatment with PEDNF

A. Treatment of Tumors

20 PEDNF is administered, alone or in conjunction with other clinical (such as surgical) procedures. PEDNF is administered by intravitreal, subretinal, intravenous or intramuscular injection or injected or applied at sites of tumor growth. The PEDNF may be administered alone or in conjunction with compounds, such as polylactic acid, which facilitate a slow, "time release" of the PEDNF. Typically, about 0.01 to about 10 μ g of PEDNF at a concentration of about 1 to about 100 μ g/ml in a physiologic saline solution or other buffered solution is administered per dose, and about 0 to about 10 doses are administered per day. When time-release compounds are included in the composition, they are used at a concentration of about 1 to about 100 μ g/ml. Treatment is continued until cessation of tumor growth and/or tumor regression is observed.

30 Treatment with PEDNF is effective for retinal tumors such as retinoblastoma, other neuronal tumors such as neuroblastoma, or tumors of non-neuronal origin. The treatment results in a cessation or reduction in the rate of cell division and a concomitant reduction in the rate of tumor growth, which in turn results in tumor regression.

B. Neurotrophic Treatment of Ocular Disease

35 PEDNF is administered, alone or in conjunction with other clinical (such as surgical) procedures. PEDNF is administered by intravitreal or subretinal injection. The PEDNF may

-21-

1 be administered alone or in conjunction with compounds such as polylactic acid which
facilitate a slow, "time release" of the PEDNF. Typically, about 0.01 to about 10 μg of
PEDNF at a concentration of about 1 to about 100 $\mu\text{g}/\text{ml}$ in a physiologic saline solution or
other buffered solution is administered per dose, and about 0 to about 10 doses are
5 administered per day. When time-release compounds are included in the composition, they
are used at a concentration of about 1 to about 100 $\mu\text{g}/\text{ml}$. Treatment is continued until the
progress of the pathology is halted and/or reversed.

Treatment with PEDNF is directed at diseases of the neural retina, retinal pigmented
epithelium, and other ocular tissue. Treatment results in enhanced survival and well-being
10 of the photoreceptors and other ocular cells, prolonging their functional life span and delaying
the onset of impaired vision and ultimate blindness.

C. Neurotrophic Treatment of Injured Nerves

PEDNF is administered, alone or in conjunction with other clinical (such as surgical)
15 procedures. PEDNF is administered by intravenous or intramuscular injection or application
or injection at the site of nerve injury. The PEDNF may be administered alone or in
conjunction with compounds such as polylactic acid which facilitate a slow, "time release"
of the PEDNF. Typically, about 0.01 to about 10 μg of PEDNF at a concentration of about
1 to about 100 $\mu\text{g}/\text{ml}$ in a physiologic saline solution or other buffered solution is
20 administered per dose, and about 0 to about 10 doses are administered per day. When time-
release compounds are included in the composition, they are used at a concentration of about
1 to about 100 $\mu\text{g}/\text{ml}$. Treatment is continued until nerve regeneration is completed.

Treatment with PEDNF is directed at injuries to nerves. Treatment results in
promotion of neurite outgrowth and nerve regeneration.

D. Treatment of Conditions Related to Serine Proteinases

PEDNF is a serine proteinase inhibitor. As such, it is effective in the treatment of
30 conditions caused by serine proteinases or where a serine proteinase inhibitor would be
advantageous. Serine proteinases include, but are not limited to, chymotrypsin, trypsin,
subtilisin, elastin, thrombin, and plasmin. Therefore, PEDNF is useful as: an anti-coagulant,
an anti-thrombotic, an anti-microbial, an anti-fungal, an anti-parasitic, and a contraceptive;
in cosmetic preparations as a proteinase inhibitor; as a weight-gain promoter; in treatments
35 for elastosis, vascular disorders involving fibrinoid formation, coagulation disorders,
arteriosclerosis, ischemia, arthroses diabetes, emphysema, arthritis, septic shock, lung
diseases, excessive complement activation, ulcers, ulcerative colitis, pancreatitis, psoriasis,

-22-

1 fibrinolytic disease, arthropathy, bone resorption, hypertension, congestive heart failure, cirrhosis, or allergy caused by proteases.

For use as an anti-coagulant, an anti-thrombotic, an anti-microbial, an anti-parasitic, or a contraceptive, or in treatment of elastosis, vascular disorders involving fibrinoid
5 formation, coagulation disorders, arteriosclerosis, ischemia, arthroses diabetes, emphysema, arthritis, septic shock, lung diseases, excessive complement activation, pancreatitis, psoriasis, fibrinolytic disease, arthropathy, bone resorption, hypertension, congestive heart failure, cirrhosis, or allergy caused by proteases, PEDNF is administered by intravenous or intramuscular injection or site-directed injection or application. The PEDNF may be
10 administered alone or in conjunction with compounds such as polylactic acid which facilitate a slow "time release" of the PEDNF. Typically, about 0.01 to about 10 μg of PEDNF at a concentration of about 1 to about 100 $\mu\text{g}/\text{ml}$ in a physiologic saline solution or other buffered solution are administered per dose, and about 0 to about 10 doses are administered per day. When time-release compounds are included in the composition, they are used at a
15 concentration of about 1 to about 100 $\mu\text{g}/\text{ml}$. Treatment is continued until progress of the pathology is halted and/or reversed.

Treatment with PEDNF is directed at injuries to nerves. Treatment results in promotion of neurite outgrowth and nerve regeneration.

For use in cosmetic preparations as a proteinase inhibitor, arthritis or elastosis PEDNF
20 is added to the preparations to a concentration of about 0.01 to about 100 $\mu\text{g}/\text{ml}$.

For use in treatment as a weight-gain promoter, ulcers, ulcerative colitis, or pancreatitis, PEDNF may be administered orally at a concentration of about 0.01 to about 10 μg per kg of body weight per day.

For use as a treatment for conditions such as psoriasis, PEDNF may be administered
25 topically at a concentration of about 0.01 to about 100 $\mu\text{m}/\text{ml}$, formulated in a suitable carrier.

Example 1

Establishment of RPE Cell Cultures

30 RPE cells were harvested from post-mortem human eyes, as described by Pfeffer (1991). The cells were grown in 75 cm flasks with Eagles's minimal essential medium supplemented with 15% fetal calf serum and 5% carbon dioxide at 37°C. Cells were grown to confluence, harvested by trypsinizing the cell monolayer, and resuspended in MEM supplemented with 15% fetal calf serum. A portion of the cells (about 5%) were reseeded
35 into new 75 cm, where the cells were again grown in 5% CO_2 at 37°C.

1

Example 2Preparation of RPE-conditioned Medium

5

Confluent cultures of RPE cells were washed extensively with HBSS, before conditioning, to remove serum proteins. Twenty-five ml of serum-free MEM was added, and the cultures were incubated with 5% CO₂ at 37°C for about 48 hours. The conditioned medium was then collected and centrifuged at 1,000 rpm at room temperature, to remove any free cells and other particulate matter, to provide an impure PEDNF protein fraction.

10

Example 3Purification of PEDNF by SDS Gel Electrophoresis

Proteins contained in human fetal RPE-conditioned medium, prepared as described in Example 2, and in control non-conditioned medium were analyzed on an SDS-polyacrylamide slab gel.

15

The conditioned and non-conditioned media samples were mixed with an electrophoresis sample buffer (62.5 mM Tris, pH 6.8, 2% SDS, 10% glycerol, 0.001% bromo- phenol blue, and 0.1 M 2-mercaptoethanol). Molecular-weight markers, such as phosphorylase B, bovine serum albumin, ovalbumin, carbonic anhydrase, soybean trypsin inhibitor and lysozyme (such as that supplied by Bio-Rad), were also mixed with an electrophoresis sample buffer. All the samples were denatured at 100°C, in a boiling water bath, for 5 min. and loaded onto a SDS containing 7.5% polyacrylamide gel. The electrophoresis was conducted at 20 mA until the bromphenol blue marker dye migrated to the bottom of the gel.

20

25

At the completion of electrophoresis, strips from each outside edge of the gels, which included the markers and a lane each of conditioned and non-conditioned media, were stained with Coomassie blue. The stained protein bands which developed on the stained strips were used to align with the unstained portion of the gel, to locate the proteins present in conditioned medium but absent from non-conditioned medium.

30

The about 50,000 to about 55,000 molecular-weight PEDNF protein doublet, unique to RPE-CM, were excised, and the proteins were electroeluted from the unstained portion of the gel. A strip of the same gel excised from the region containing bovine serum albumin was treated similarly to serve as a control. The eluant was centrifuged to remove gel fragments, and the supernatant was dialyzed and analyzed by gel electrophoresis to assess its purity. The electrophoretic pattern of the SDS-polyacrylamide gel electrophoresis is shown in FIG. 1.

35

The electrophoretic patterns of human fetal RPE-CM and non-conditioned control medium show the presence of the prominent about 50,000 to about 55,000 molecular-weight PEDNF doublet unique to the conditioned medium.

-24-

Example 4Small Scale Ammonium Sulfate Precipitation

An impure PEDNF protein fraction, prepared as described in Example 2, was divided into six 10 ml aliquots. One of the aliquots was brought to 40% saturation, at 20°C, by the addition of 2.42 g of ammonium sulfate; another aliquot was brought to 50% saturation, at 20°C, by the addition of 3.14 g of ammonium sulfate; another aliquot was brought to 60% saturation, at 20°C, by the addition of 3.90 g of ammonium sulfate; another aliquot was brought to 70% saturation, at 20°C, by the addition of 4.72 g of ammonium sulfate; another aliquot was brought to 80% saturation, at 20°C, by the addition of 5.61 g of ammonium sulfate; and the final aliquot was brought to 90% saturation, at 20°C, by the addition of 6.57 g of ammonium sulfate. The aliquots were mixed until the ammonium sulfate was completely dissolved. The precipitates which formed in each of the tubes were collected by centrifugation at 10,000 rpm for 20 min. The precipitates were each resuspended in 10 mM sodium phosphate buffer, pH 7.5, and dialyzed against 2 l of 10 mM sodium phosphate buffer, pH 7.5, for 24 hours.

The samples were then collected and subjected to SDS-polyacrylamide gel electrophoresis on a 10% SDS-polyacrylamide gel, as described in Example 3.

The results are summarized in Table I.

Table I

<u>% Saturation Ammonium Sulfate</u>	<u>Relative concentration of PEDNF</u>
40	-
50	-
60	+
70	+++
80	-
90	-

The results indicate that the PEDNF precipitates with an ammonium sulfate saturation of about 60% to 70%. The small amount of PEDNF in the 50% to 60% sample indicates that some PEDNF is present, and that a suitable ammonium sulfate "cut" is from 50% to 70% saturation. The samples were subjected to SDS-polyacrylamide gel electrophoresis, as described in Example 3. It was estimated that the purity of the PEDNF in the 60% to 70% ammonium sulfate fraction was about 50%.

-25-

1

Example 5Ammonium Sulfate Precipitation

5

An impure protein fraction, prepared as described in Example 2, was brought to 50% saturation with ammonium sulfate by the addition of 313 g/l of solid ammonium sulfate. The ammonium sulfate was added slowly to the impure protein fraction while the solution was stirred with a stir bar on a mechanical stirrer. After all the ammonium sulfate was added, the 50% ammonium sulfate solution was stirred for 30 min. at 20°C. The 50% ammonium sulfate solution was then centrifuged at 10,000 g, at 20°C for 20 min. The supernatant was collected.

10

The 50% ammonium sulfate solution was then brought to 70% saturation by the addition of 137 g/l of ammonium sulfate. The ammonium sulfate was added slowly, and after all the ammonium sulfate was added, the 70% ammonium sulfate solution was stirred for 30 min. at 20°C. The 70% ammonium sulfate solution was centrifuged or filtered, as described above, and the precipitate was collected.

15

The ammonium sulfate precipitate was then redissolved in 10 mM sodium phosphate, pH 7, to form a 55%-70% ammonium sulphate fraction, then diafiltered using an Amicon Diaflo ultrafiltration unit, or dialyzed against 10 mM sodium phosphate, pH 7, to remove the ammonium sulfate from the redissolved 55-70% fraction.

20

The samples were subjected to SDS-polyacrylamide gel electrophoresis, as described in Example 3. It was estimated that the purity of the PEDNF in the 55% to 70% ammonium sulfate fraction was about 50%.

Example 6Purification of PEDNF by Cation Exchange HPLC

25

RPE-CM, prepared as described in Example 2, was dialyzed against 10 mM sodium phosphate buffer, pH 6.5, to remove media and salts from the impure protein sample. The dialyzed PEDNF sample was then loaded onto a Brownlee Aquapore CX-300 HPLC chromatography medium, packed into a 4.6 X 30 mm column. The chromatography medium was previously equilibrated with 10 mM phosphate, pH 6.5. The dialyzed RPE-CM was loaded onto the chromatography medium, and the chromatography medium, with PEDNF, was washed with 10 mM phosphate, pH 6.5, until all unbound proteins were washed from the chromatography medium. PEDNF was eluted from the chromatography medium with a linear salt gradient from 0.0 to about 0.5 M NaCl in 10 mM phosphate, pH 6.5. PEDNF, eluted with a NaCl concentration of about 0.25 M. The eluted PEDNF was concentrated by lyophilization and resolubilized in 10 mM phosphate, pH 6.5. The eluted PEDNF was analyzed by SDS gel electrophoresis, as described in Example 3. The eluted PEDNF was estimated to be approximately 70% pure.

30

35

1

Example 7Purification of PEDNF by Cation-Exchange HPLC

5

Ammonium sulfate-precipitated RPE-CM, prepared as described in Example 5, was dialyzed against 10 mM sodium phosphate buffer, pH 6.5, to remove ammonium sulfate. An 80 μ l sample of the dialyzed ammonium sulfate-purified PEDNF protein fraction was loaded onto a Brownlee Aquapore CX-300 chromatography medium, packed into a 4.6 X 30 mm column. The chromatography medium was previously equilibrated with 10 mM phosphate, pH 7.2. The PEDNF/chromatography medium was washed with 10 mM sodium phosphate buffer, pH 7.2, until all unbound proteins were washed from the chromatography medium. PEDNF was eluted from the chromatography medium with a linear salt gradient from 0.0 to about 0.5 M NaCl in 10 mM sodium phosphate buffer, pH 7.2. PEDNF eluted as a single peak with a NaCl concentration of about 0.25 M.

10

15

The eluted PEDNF was dialyzed against 10 mM sodium phosphate buffer, pH 7.2, lyophilized, and resolubilized in water. The sample was then subjected to SDS-polyacrylamide gel electrophoresis, as described in Example 3.

The eluted PEDNF was estimated to be about 70% pure.

Example 8Purification of PEDNF by Reverse-Phase HPLC

20

Ammonium sulfate-precipitated RPE-CM, prepared as described in Example 5, was dialyzed against 10 mM sodium phosphate buffer, pH 6.5, to remove ammonium sulfate. An 80 μ l PEDNF-containing sample was then loaded onto a Vydac C8 chromatography medium, packed into a 4.6 X 250 mm column. The chromatography medium was previously equilibrated with 0.1% TFA in water. The PEDNF/chromatography medium was washed with 0.1% TFA in water until all unbound proteins were washed from the chromatography medium. PEDNF was eluted from the chromatography medium with a linear gradient from 0.0 to about 95% CH₃CN, 0.1% TFA in water, and 5% H₂O.

25

30

The eluted PEDNF was dialyzed against 10 mM sodium phosphate buffer, pH 7.2, lyophilized, and resolubilized in water. The sample was then subjected to SDS-polyacrylamide gel electrophoresis, as described in Example 3.

The PEDNF eluted from the column was estimated to be about 80% pure.

Example 9Purification of PEDNF by Reverse-Phase HPLC

35

A PEDNF-containing sample, partially purified as described in Example 2, was then loaded onto a Vydac C8 chromatography medium, packed into a 4.6 X 250 mm column. The chromatography medium was previously equilibrated with 0.1% TFA in water. The

-27-

1 PEDNF/chromatography medium was washed with 0.1% TFA in water until all unbound proteins were washed from the chromatography medium. PEDNF was eluted from the chromatography medium with a linear gradient from 0.0 to about 95% CH₃CN in 0.1% TFA and 5% H₂O.

5 The eluted PEDNF was dialyzed against 10 mM sodium phosphate buffer, pH 7.2, lyophilized, and resolubilized in water. The sample was then subjected to SDS-polyacrylamide gel electrophoresis, as described in Example 3.

The PEDNF eluted from the column was estimated to be about 60% pure.

10

Example 10

Purification of PEDNF by Reverse-Phase HPLC

A PEDNF-containing sample, partially purified as described in Example 6, was then loaded onto a Vydac C8 chromatography medium, packed into a 4.6 X 250 mm column. The chromatography medium was previously equilibrated with 0.1% TFA in water. The PEDNF /chromatography medium was washed with 0.1% TFA in water until all unbound proteins were washed from the chromatography medium. PEDNF was eluted from the chromatography medium with a linear gradient from 0.0 to about 95% CH₃CN in 0.1% TFA and 5% H₂O.

20 The eluted PEDNF was dialyzed against 10 mM sodium phosphate buffer, pH 7.2, lyophilized, and resolubilized in water. The sample was then subjected to SDS-polyacrylamide gel electrophoresis, as described in Example 3.

The PEDNF eluted from the column was estimated to be about 80% pure.

25

Example 11

Purification of PEDNF by Reverse-Phase HPLC

A PEDNF-containing sample, prepared as described in Example 6, was loaded onto Brownlee RP-300 chromatography medium, packed into a 4.6 X 250 mm column. The chromatography medium was previously equilibrated with 0.1% TFA in water. The PEDNF /chromatography medium was washed with 0.1% TFA in water until all unbound proteins were washed from the chromatography medium. PEDNF was eluted from the chromatography medium with a linear gradient from 0.0 to about 95% CH₃CN in 0.1% TFA and 5% H₂O.

30 The eluted PEDNF was dialyzed against 10 mM sodium phosphate buffer, pH 7.2, lyophilized, and resolubilized in water. The sample was then subjected to SDS-polyacrylamide gel electrophoresis, as described in Example 3.

The PEDNF eluted from the column was estimated to be about 90% pure.

Example 12Purification of PEDNF by Size-Exclusion Chromatography

Partially-purified PEDNF, prepared as described in Example 5 or 6, was purified further by chromatography on Bio-Rad TSK-250 chromatography medium, packed into a 7.5 X 300 mm column. A 40 μ l sample of resolubilized PEDNF was loaded onto the size-exclusion chromatography medium, which had been previously equilibrated with 20 mM Tris, pH 7.0, 0.6 M NaCl. PEDNF was eluted with 20 mM Tris, pH 7.0, 0.6 M NaCl.

The eluted PEDNF was dialyzed against 10 mM sodium phosphate buffer, pH 7.2, lyophilized, and resolubilized in water. The sample was then subjected to SDS-polyacrylamide gel electrophoresis, as described in Example 3.

The PEDNF eluted from the column was estimated to be about 75% pure.

Example 13Anion-Exchange Chromatography

DEAE cellulose is equilibrated with 10 mM Tris, pH 7.5. PEDNF, prepared as described in Example 5, is applied to the column, and the column is washed with 10 mM Tris, pH 7.5, until all unbound proteins are eluted from the column. After all the unbound proteins are eluted, the PEDNF is eluted with a linear gradient from 0 to about 1 M NaCl in 10 mM Tris-HCl, pH 7.5. The fractions containing PEDNF are collected and pooled. These fractions are then diafiltered against 10 mM Tris-HCl, pH 7.5 to remove NaCl.

Example 14Anion-Exchange Chromatography

DEAE cellulose is equilibrated with 10 mM Tris, pH 7.5. PEDNF, prepared as described in Example 2, is applied to the column, and the column is washed with 10 mM Tris, pH 7.5, until all unbound proteins are eluted from the column. After all the unbound proteins are eluted, the PEDNF is eluted with a linear gradient from 0 to about 1 M NaCl in 10 mM Tris-HCl, pH 7.5. The fractions containing PEDNF are collected and pooled. These fractions are then diafiltered against 10 mM Tris-HCl, pH 7.5 to remove NaCl.

Example 15Cation-Exchange Chromatography

Bio-Rex 70 resin (Bio-Rad) cellulose is equilibrated with 10 mM PBS, pH 7.2. PEDNF, prepared as described in Example 2, is applied to the column, and the column is washed with 10 mM PBS, pH 7.2, until all unbound proteins are eluted from the column. After all the unbound proteins are eluted, the PEDNF is eluted with a linear gradient from 0 to about 1 M NaCl in 10 mM PBS, pH 7.2. The fractions containing PEDNF are collected

1 and pooled. These fractions are then diafiltered against 10 mM Tris-HCl, pH 7.5 to remove NaCl.

Example 16

Cation-Exchange Chromatography

5 Bio-Rex 70 resin (Bio-Rad) cellulose is equilibrated with 10 mM PBS, pH 7.2. PEDNF, prepared as described in Example 5, is applied to the column, and the column is washed with 10 mM PBS, pH 7.2, until all unbound proteins are eluted from the column. After all the unbound proteins are eluted, the PEDNF is eluted with a linear gradient from 0 to about 1 M NaCl in 10 mM PBS, pH 7.2. The fractions containing PEDNF are collected
10 and pooled. These fractions are then diafiltered against 10 mM Tris-HCl, pH 7.5 to remove NaCl.

Example 17

Heparin Chromatography

15 Two ml of heparin agarose was packed into a 0.7 x 10 cm column, washed with 20 ml of 10 mM Tris-HCl, pH 7.5, 3 M NaCl, then equilibrated with 20 ml of 10 mM Tris-HCl, pH 7.5. Twelve ml of a PEDNF solution, prepared as described in Example 2, from a 17-year-old human donor, was dialyzed against 4 l of 10 mM sodium phosphate buffer, pH 7.5, at 4°C for 19 hours. The volume after dialysis was 19 ml. The dialysate was applied
20 to the heparin agarose at a flow rate of 0.4 ml/min. After the PEDNF solution had been applied to the heparin agarose, the heparin agarose was washed with 20 ml of 10 mM Tris-HCl, pH 7.5, to remove unbound proteins from the heparin agarose. PEDNF was then eluted from the heparin agarose with 12 ml of 10 mM Tris-HCl, pH 7.5, 3 M NaCl.

The eluate containing PEDNF was dialyzed against 10 mM Tris-HCl, pH 7.5, to
25 remove the NaCl present in the eluate. The dialyzed eluate was then lyophilized, redissolved in 10 mM sodium phosphate buffer, pH 7.5, and a sample was subjected to 12.5% SDS-polyacrylamide gel electrophoresis, as described in Example 3.

The PEDNF eluate was estimated to be about 60% pure.

30

Example 18

Heparin Chromatography

Two ml of heparin agarose was packed into a 0.7 x 10 cm column, washed with 20 ml of 10 mM Tris-HCl, pH 7.5, 3 M NaCl, then equilibrated with 20 ml of 10 mM Tris-HCl, pH 7.5. One ml of a PEDNF solution, prepared as described in Example 5, was
35 dialyzed against 4 l of 10 mM sodium phosphate buffer, pH 7.5, at 4°C for 19 hours. The volume of the dialysate was 1.3 ml. The dialysate was applied to the heparin agarose at a flow rate of 0.4 ml/min. After the PEDNF solution had been applied to the heparin agarose,

-30-

1 the heparin agarose was washed with 20 ml of 10 mM Tris-HCl, pH 7.5, to remove unbound
proteins from the heparin agarose. The heparin agarose was then successively eluted with
12 ml of 10 mM Tris-HCl, pH 7.5, 0.5 M NaCl; 12 ml of 10 mM Tris-HCl, pH 7.5, 1 M
NaCl; 12 ml of 10 mM Tris-HCl, pH 7.5, 2 M NaCl; and 12 ml of 10 mM Tris-HCl, pH
5 7.5, 3 M NaCl.

Each of the eluates was dialyzed against 10 mM Tris-HCl, pH 7.5, to remove the
NaCl present in the eluate. The dialyzed eluates were then lyophilized and redissolved in 10
mM sodium phosphate buffer, pH 7.5, and a sample of each dialyzed eluate was subjected
to 12.5% SDS-polyacrylamide gel electrophoresis, as described in Example 3.

10 The PEDNF eluate was estimated to be about 70% pure.

Example 19

Neuronal Inductivity and Differentiation Activity of Isolated PEDNF

15 Electroeluted PEDNF was prepared as described in Example 3. To characterize the
neuronal inductive activity of the isolated PEDNF, the optimal concentration and pre-
seeding stimulatory periods were determined. Either 1, 2, 4, 6, 8, or 10 $\mu\text{g/ml}$ of
electroeluted PEDNF in serum-free DME, supplemented with 1 mM sodium pyruvate, 0.625
mM HEPES, 6 mM L-glutamine, 1% non-essential amino acids, 5 $\mu\text{g/ml}$ insulin, 5 $\mu\text{g/ml}$
20 transferrin, and 5 ng/ml selenous acid, was tested. As a control, RPE-CM, diluted with an
equal volume of non-conditioned medium (50% RPE-CM), was also used.

The Y79 RB cells were grown in suspension culture. The cells were harvested by
centrifugation for 5 min. at 900 rpm at room temperature and resuspended in serum-free
DME medium, which had previously been warmed to 37°C. The cells were collected by
25 centrifugation and resuspended at a concentration of 10^6 cells/ml in serum-free DME medium.
One, 2, 4, 6, 8, or 10 $\mu\text{g/ml}$ electro-eluted PEDNF or 50% RPE-CM was introduced into
separate Y79 RB cell cultures. The cells were incubated for 7 days at 37°C, and were then
attached to poly-D-lysine-coated flasks. The cells were observed daily by phase-contrast
microscopy.

30 The optimal pre-seeding period required for maximal inductive activity of PEDNF was
assessed by treating Y79 RB cell cultures with 2 $\mu\text{g/ml}$ of electroeluted PEDNF for 2, 4, 8,
or 16 days prior to seeding onto a poly-D-lysine substratum. In addition, 1 or 2 $\mu\text{g/ml}$ of
electroeluted PEDNF was added to attached cultures of cells not previously stimulated in
suspension culture. Greater than 80% of Y79 cell aggregates treated with 2 $\mu\text{g/ml}$
35 electroeluted PEDNF extended arborizing processes within 8-10 days after attachment to a

-31-

poly-D-lysine substratum; untreated cells showed only minimal signs of neuronal differentiation (FIG. 2).

Y79 cells exhibited maximal differentiative response to electroeluted PEDNF at a protein concentration of 2 $\mu\text{g/ml}$; the response was reduced at concentrations exceeding 4 $\mu\text{g/ml}$. The results are shown in Table II.

Table II

	<u>PEDNF ($\mu\text{g/ml}$)</u>	<u>Pre-Attachment Stimulatory Period (days)</u>	<u>Percent Differentiated Cells</u>
10	0	7	13 \pm 8*
	1	7	82 \pm 10
	2	7	80 \pm 10
	4	7	33 \pm 10
	6	7	5 \pm 5
15	8	7	0
	10	7	0
	1	0 ⁺	20 \pm 14
	2	0 ⁺	28 \pm 11
20	1 $\mu\text{g/ml}$ BSA (control)	7	8 \pm 6

* Standard error

⁺ PEDNF added to non-stimulated attached cells,
24 hrs. post-seeding.

A pre-seeding stimulatory period (in suspension culture) of 8 days, in the presence of this factor, results in maximal differentiation of Y79 cells at day 10 post-attachment. Decreased frequencies of differentiation (less than 50%) are noted with both shorter and longer pre-seeding inductive periods. In addition, cells not previously stimulated with PEDNF prior to attachment, exhibit a low frequency of neuronal differentiation (approximately 20%) if 1-2 $\mu\text{g/ml}$ electroeluted protein is added post-attachment. This response, however, is relatively slow, i.e., less than 15 days. Control BSA-treated cultures exhibit less than 10% differentiation, comparable to cells exposed only to non-conditioned media.

PEDNF induced a high degree of neuronal differentiation in human retinoblastoma cells. Long ramifying neuritic processes were induced to extend from aggregates of stimulated cells; concomitant increases in the expression of the neuronal marker molecules, neuron-specific enolase, and the 200,000 molecular weight neurofilament protein were also

observed. The expression of the neuronal phenotype in Y79 cells appears to involve three sequential events: 1) stimulation of cells in suspension culture by PEDNF; 2) attachment of stimulated cells to a substratum; followed by 3) neurite outgrowth of attached, stimulated cells. Since only 20% differentiation was observed when non-stimulated, attached cultures were treated with PEDNF, commitment to neuronal differentiation in Y79 cells appears to precede cell attachment and neurite outgrowth and may be initiated during the stimulatory period.

Example 20

Comparison of the Effects of RPE-CM from Different Species

Cell cultures were established as described in Example 1, except the eyes from which the cells were derived were either fetal humans, adult humans, adult rats, fetal rats, or embryonic chickens.

RPE-CM was prepared from each of the cell cultures, as described in Example 2. Fetal human cell cultures which were newly established (short-term cultures) and cultures which had been established for 6 months (long-term cultures) were included.

The differentiation activity of the conditioned media was performed as described in Example 19.

The results of experiments monitoring the effects of RPE-CM from various species are summarized in Table III.

Table III

<u>Source of RPE-Conditioned Medium Aggregates</u>	<u>% Differentiated</u>
Human (fetal, short-term cultures)	88 \pm 3*
Human (fetal, long-term cultures)	50 \pm 8
Human (adult)	55 \pm 9
Rat (adult)	20 \pm 11
Rat (fetal)	42 \pm 7
Chicken (embryonic)	86 \pm 7

* Standard error

The table summarizes the inductive activity of a variety of RPE-conditioned media on Y79 cells. Indicated are the percentages of 10 days attached aggregates (more than 5 cells/aggregate) exhibiting neurite outgrowth after 7 days' stimulation with 50% RPE-CM from the various species. One hundred aggregates were counted from each sample in replicates of three.

-33-

1 The greatest differentiative response in Y79 cells is induced by human fetal RPE-CM
(short-term cultures) and embryonic chicken RPE-CM, both of which induce neuronal
differentiation in greater than 80% of cellular aggregates when used at a 50% concentration.
In contrast, decreased frequencies (less than 50%) of cellular differentiation are noted for
5 conditioned media from long-term (12-18 months) cultures of human fetal RPE and adult
human RPE. Only 40% neuron-like differentiation is induced by fetal rat RPE-CM and 20%
by adult rat RPE-CM.

 Conditioned media from human fetal and embryonic chicken RPE cells contain similar
neurotrophic activity, while those from adult cultures (chicken and rat) and long-term cultures
10 of human fetal RPE-CM are less effective in promoting the neuronal phenotype in Y79 cells.
It is possible that mature RPE cells no longer secrete this neurotrophic factor, or they secrete
reduced quantities which are less effective in the culture conditions utilized. The fact that
some trophic activity is seen in RPE-conditioned media from other species suggests that
PEDNF, or a similar factor, is also secreted by RPE cells of other species and may be
15 generally important to normal retinal development and function.

Example 21

Two-Dimensional Gel Electrophoresis

 Two-dimensional gel analysis was conducted by the method described by O'Farrell,
20 J. Biol. Chem., **250**, 4007-4021 (1975) and Jones, J., J. Exp. Med., **14b**, 1251-1279 (1977).
Silver-staining was performed using a Bio-Rad silver-stain plus kit. Twenty μ g of PEDNF,
purified by cation-exchange and size-exclusion HPLC, was loaded onto the IEF tube gel.

 Two-dimensional electrophoretic analysis of HPLC-purified PEDNF reveals the
presence of four closely-grouped molecular species (FIG. 3). The four species vary slightly
25 in apparent molecular weight, but all are in the 50,000 molecular-weight region of the gel,
consistent with the previously identified molecular weight of PEDNF. The resolved species
also vary slightly in isoelectric point, suggesting slight variations in post-translational
modifications. Amino-acid sequence analysis (see Example 22), however, indicated the
presence of only a single polypeptide.

Example 22

Isolation of PEDNF Specific Peptides and Amino-Acid Sequences

 180 μ g of purified PEDNF (purified as described in Examples 2, 6, and 12) was
35 concentrated using a Centricon 10 microconcentrator, supplied by Amicon of Danvers, MA,
and then diluted in 25 mM Tris, pH 8.5, 1 mM EDTA. To the protein sample was added

-34-

1 endoproteinase Lys-C. The PEDNF/proteinase mixture was incubated for about 18 hrs. at 30°C to digest the PEDNF.

The resulting PEDNF polypeptide fragments were separated using by HPLC on a Vydac C8 reverse-phase HPLC packed into a 4.6 X 250 mm column. The column was
5 equilibrated with 0.1% TFA in water. The polypeptides were eluted with 90% CH₃CN, 0.1% TFA in water.

Polypeptides eluted from the column which are well separated from other polypeptides were collected and subjected to protein sequencing analysis. The amino-acid sequence analysis was performed under contract at the Microsequencing Facility of Beckman Research
10 Institute at the City of Hope.

The amino-acid sequences for the isolated polypeptides were determined to be:

PEDNF-13 (ID SEQ NO 1 residues 226-244)

Thr Ser Leu Glu Asp Phe Tyr Leu Asp Glu Glu Arg Thr Val Arg Val Pro Met Met

15 PEDNF-14 (ID SEQ NO 1 residues 161-186)

Ser Tyr Gly Thr Arg Pro Arg Val Leu Thr Gly Asn Pro Arg Leu Asp Leu Gln Glu Ile Asn
Asn Trp Val Gln Ala

Sequencing of PEDNF-13 yielded 19 residues of unequivocal sequence. PEDNF-14 yielded 26 residues, 25 of which were unequivocal. The identification of residue 23 in
20 PEDNF-14 as tryptophan was not absolutely certain.

Example 23

Isolation of PEDNF Specific Peptides and Amino Acid Sequence

25 PEDNF, purified as described in Examples 2, 6, and 12, was reduced and alkylated. The sample was dried, redissolved in 50 µl of CRA buffer (8 M urea, 0.4 M ammonium bicarbonate, pH 8.0), and 5 µl of 45 mM DTT (Calbiochem) was added. After heating at 50°C for 15 minutes, the solution was cooled, and 5 µl of 100 mM iodoacetic acid (Sigma Chemical Co.) was added. After 15 minutes, the solution was diluted to a concentration of
30 2 M urea and subjected to trypsin digestion, supplied by Boehringer-Mannheim, using an enzyme:substrate ratio of 1:25 (wt/wt), for 22 hrs at 37°C.

Tryptic peptides were separated by narrow-bore reverse-phase HPLC on a Hewlett-Packard 1090 HPLC, equipped with a 1040 diode array detector, using a Vydac 2.1 mm x 150 mm C18 column. Buffer A was 0.06% trifluoroacetic acid/H₂O, and buffer B was
35 0.055% trifluoroacetic acid/acetonitrile, a gradient of 5% B at 0 min., 33% B at 63 min., 60% B at 95 min., and 80% B at 105 min., with a flow rate of 150 µl/min., was used. Chromatographic data at 210, 277 nm and UV spectra from 209 to 321 nm of each peak were

-35-

1 obtained. Samples for amino terminal sequence analysis were applied to a polybrene pre-
cycled glass fibre filter and subjected to automated Edman degradation at the Harvard
Microchemical Facility in Boston, MA, on an ABI model 477A gas-phase protein sequencer
using program NORMAL 1. The resulting phenylthiohydantoin amino acid fractions were
5 manually identified using an on-line ABI Model 120A HPLC and Shimadzu CR4A integrator.

The sequences of the isolated peptides were determined to be:

PEDNF 1 (ID SEQ NO. 1 residues 54-67)

Leu Ala Ala Ala Val Ser Asn Phe Gly Tyr Asp Leu Tyr Arg

10 PEDNF 2 (ID SEQ NO. 1 residues 107-135)

Ala Leu Tyr Tyr Asp Leu Ile Ser Ser Pro Asp Ile His Gly Thr Tyr Lys Glu Leu Leu Asp Thr
Val Thr Ala Pro Gln Lys Asn

PEDNF 5 (ID SEQ NO. 1 residues 307-316)

Thr Val Gln Ala Val Leu Thr Val Pro Lys

15 PEDNF 6 (ID SEQ NO. 1 residues 317-327)

Leu Lys Leu Ser Tyr Glu Gly Glu Val Thr Lys

PEDNF 7 (ID SEQ NO. 1 residues 360-389)

Ala Gly Phe Glu Trp Asn Glu Asp Gly Ala Gly Thr Thr Pro Ser Pro Gly Leu Gln Pro Ala
His Leu Thr Phe Pro Leu Asp Tyr His

20 Example 24

Comparison of the PEDNF Peptides
to Rat Serine Proteinase Sequences

PEDNF-13 shows significant sequence. Homologous molecules include rat serine
protease inhibitors 1 and 2, a rat hepatocyte growth hormone-regulated protein, a rat thyroid
25 hormone-regulated protein, and mouse contrapsin. Human, monkey, sheep, and mouse alpha-
1 antitrypsin, and porcine alpha-1-antichymotrypsin, show significant but somewhat less
homology. PEDNF-14 shows homology with different protease inhibitors, but the degree of
homology is less than that observed for PEDNF-13. These include human plasma protease
C1 inhibitor and human alpha-2-antiplasmin inhibitor. These results suggest that PEDNF may
30 function as a protease inhibitor, but that it is molecularly distinct from such inhibitors which
have been described previously.

PEDNF shows significant homology with serpins, the family of serine protease
inhibitors that share a common reactive center near the C-terminal end, which serves as an
exposed binding site that acts as bait for target serine proteinases. It is of interest that a
35 number of known members of the serpin family have been shown to have neurotrophic effects
on a variety of neuronal cell types. For example, glial-derived nexin (GDN) promotes neurite
outgrowth in neuroblastoma cells, as do a number of other protease inhibitors, such as hirudin

-36-

1 and leupeptin. Protease inhibitors also stimulate neuronal differentiation in cells of dorsal
root ganglia, sympathetic neurons, and hippocampal pyramidal neurons. It is also of interest
that the production of proteinases and protease inhibitors is stimulated by a number of known
growth factors. While it remains to be determined if PEDNF has protease inhibitor activity,
5 it is likely, since inhibitory activity has been shown to be necessary for the neurite-promoting
activity of glial derived nexin. It has been suggested that the formation of a stable protease-
GDN complex, and a consequent conformational change in GDN and/or the associated
protease, is necessary for the neurite-promoting activity of GDN. PEDNF may well function
similarly, as there are known proteinases present in the interphotoreceptor matrix and in the
10 developing neural retina.

Example 25

Cloning of the PEDNF Gene

Oligonucleotides were constructed against PEDNF 13:

15 5'-AGYAAATTYTAYGAYCTSTA-3'

determined in Example 22, and PEDNF 2:

5'-CTYTCYTCRTCSAGRTARAA-3'

determined in Example 23.

The oligonucleotides were prepared on an ABI 392 DNA/RNA Synthesizer and used
20 as primers in a polymerase chain reaction (PCR). A human fetal eye Charon BS cDNA
library, donated by Dr. A. Swaroop, was amplified once by the method described by
Sambrook et al. (1989) Molecular Cloning: A Laboratory Manual. 2nd Ed. (Cold Spring
Harbor Press, Cold Spring Harbor, NY) and screened by PCR as described by Friedman et
al. (1990, in Screening of λ gt 11 libraries in PCR protocols: a guide to methods and
25 applications. Innis, Gelfand, Sninsky and White, eds., Academic Press, pp. 253-260) using
a Techne thermal cycler and standard reagents (GeneAMP, Perkin Elmer Cetus), except that
MgSO₄ was used at 3 mM.

The recovered fragment was isolated on a 3% NuSieve 3:1 gel (FMC Biochemicals)
using NA-45 DEAE-cellulose paper (Schleicher and Schull) and labelled with ³²P by random
30 priming (Prime-It Random Primer Labeling Kit, Stratagene).

This probe was used to screen 200,000 pfus of the same library (10). Positive clones
were isolated as described by Sambrook et al supra, and the DNA was purified with Qiagen
Maxi preparation protocols (Qiagen, Inc.).

The inserts were cut out with Not1 (BRL, Gaithersburg, MD) circularized with T4
35 DNA ligase (New England Biolabs), transformed into competent E. coli Sure cells
(Stratagene, Inc.), and plated out on 12.5 μ g/ml Ampicillin/40 μ g/ml X-gal agar plates.

-37-

1 White colonies were selected and mini-prepped (Qiagen plasmid mini-prep protocol).
Purified plasmids were digested with EcoR1/HindIII (BRL) and run on a 0.7% agarose gel
to determine the size of the inserts.

5 Example 26

DNA Sequence Analysis

One of the identified clones which contained an insert corresponding to the PEDNF
gene was selected for mapping and subsequent sequencing using a USB Sequenase 2.0
protocol and reagents.

10 Example 27

Treatment of Retinal Tumors

PEDNF is administered, alone or in conjunction with clinical (such as surgical)
procedures. PEDNF is administered by intravitreal or subretinal injection. The PEDNF may
15 be administered alone or in conjunction with compounds such as poly(lactic acid) which
facilitate a slow, "time release" of the PEDNF. Typically, about 0.01 to about 10 μg of
PEDNF at a concentration of about 1 to about 100 $\mu\text{g}/\text{ml}$ in a physiologic saline solution or
other buffered solution is administered per dose, and about 0 to about 10 doses are
administered per day. When time-release compounds are included in the composition, they
20 are used at a concentration of about 1 to about 100 $\mu\text{g}/\text{ml}$. Treatment is continued until
tumor progression is halted or reversed.

Treatment with PEDNF is effective for retinal tumors such as retinoblastoma, other
neuronal tumors such as neuroblastoma, or tumors of non-neuronal origin. The treatment
results in a cessation or reduction in the rate of cell division and a concomitant reduction in
25 the rate of tumor growth, which in turn results in tumor regression.

Example 28

Neurotrophic Treatment of Ocular Disease

PEDNF is administered, alone or in conjunction with clinical (such as surgical)
30 procedures. PEDNF is administered by intravitreal or subretinal injection. The PEDNF may
be administered alone or in conjunction with compounds such as poly(lactic acid) which
facilitate a slow, "time release" of the PEDNF. Typically, about 0.01 to about 10 μg of
PEDNF at a concentration of about 1 to about 100 $\mu\text{g}/\text{ml}$ in a physiologic saline solution or
other buffered solution is administered per dose, and about 0 to about 10 doses are
35 administered per day. When time-release compounds are included in the composition, they
are used at a concentration of about 1 to about 100 $\mu\text{g}/\text{ml}$. Treatment is continued until the
pathology is halted or reversed.

1 Treatment with PEDNF is directed at diseases of the neural retina, retinal pigmented epithelium, and other ocular tissue. Treatment results in enhanced survival and well-being of the photoreceptors and other ocular cells, prolonging their functional life span and delaying the onset of impaired vision and ultimate blindness.

5

Example 29

Neurotrophic Treatment of Injured Nerves

PEDNF is administered, alone or in conjunction with clinical (such as surgical) procedures. PEDNF is administered by intravitreal or subretinal injection. The PEDNF may be administered alone or in conjunction with compounds such as poly(lactic acid) which facilitate a slow, "time release" of the PEDNF. Typically, about 0.01 to about 10 μg of PEDNF at a concentration of about 1 to about 100 $\mu\text{g}/\text{ml}$ in a physiologic saline solution or other buffered solution is administered per dose, and about 0 to about 10 doses are administered per day. When time-release compounds are included in the composition, they are used at a concentration of about 1 to about 100 $\mu\text{g}/\text{ml}$. Treatment is continued until nerve regeneration is completed.

15

Treatment with PEDNF is directed at injuries to nerves. Treatment results in promotion of neurite outgrowth and nerve regeneration.

The above descriptions of exemplary embodiments of methods for retinal pigmented epithelium derived neurotrophic factor are for illustrative purposes. Because of variations which will be apparent to those skilled in the art, the present invention is not intended to be limited to the particular embodiments described above. The present invention may also be practiced in the absence of any element not specifically disclosed. The scope of the invention is defined by the following claims.

20

25

Example 30

Transfection of Bacterial Cells

Competent E. coli Sure cells are mixed with the ligation mixture and incubated on ice for 60 min. The mixture is then incubated at 42°C for 2 min., then 800 μl of 2% BACTO Tryptone, 2% BACTO Yeast Extract (both supplied by DIFCO LABORATORIES), 10 mM NaCl, 10 mM MgSO_4 , 20 mM glucose are added and the mixture incubated at 37°C for 60 min. 50 to 500 μl of the DNA mixtures are then spread on selection plates containing 12.5 $\mu\text{g}/\text{ml}$ Ampicillin.

30

The recombinant colonies are screened by miniprepping and digestion of the isolated DNA. Positive colonies containing the appropriate inserts are then prepared by growing the appropriate colony in 200 ml of Luria Broth plus 12.5 $\mu\text{g}/\text{ml}$ ampicillin with shaking at 37°C overnight. After the overnight incubation, the cultures are transferred to centrifuge bottles

35

-39-

1 and centrifuged at 2500 rpm for 10 min. The supernatant is removed and the cells are
resuspended in 30 ml of STET buffer (0.23 M sucrose, 5% Triton-X-100, 20 mM EDTA,
50 mM Tris-HCl, pH 8) at room temperature. Five μ l of 10 mg/ml lysozyme is added and
the mixture is swirled and incubated for about 5 min. at room temperature. Each flask is
5 then gently swirled directly over a flame until the cells begin to coagulate and turn white.
The mixture is then transferred to a boiling-water bath for about 45 sec. The solution is then
cooled in an ice-water bath for 2 min., and the mixture is transferred to centrifuge tubes and
centrifuged for 15 min. at 16,000 rpm.

The supernatant is then transferred to a clean container. Two volumes of 100%
10 ethanol are added, mixed, and the DNA precipitated at -70°C for 20 min. The precipitate
is collected by centrifugation at 2,500 g for 15 min. The ethanol supernatant is removed, and
the pellet is washed with about 10 ml of cold (-20°C) 90% ethanol. 2.5 ml of extraction
buffer (0.2 M Tris, pH 7.5, 0.08 M EDTA, and 0.2 M KCl) is added to the pellet, and it is
resuspended, at 4°C . 100 μ l of 10 mg/ml RNase, dissolved in 0.1X TE (1 mM Tris-HCl,
15 0.1 mM EDTA, pH 7.6) and pretreated by boiling for 10 min.

Example 30

Insect Cell Cultures

Sf9 cells are seeded into spinner flasks, containing Grace's Antheraea medium supplied
20 by GIBCO of Grand Island, NY, to an initial density of 1×10^6 cells/ml and incubated at
 27°C with constant stirring at 50 rpm. The cells are subcultured when they reach a density
of 2.5×10^6 cells/ml, approximately 2 to 3 times a week, and are diluted about 1 in 5.

Example 31

Cloning Genes into pAC373

Two μ l of pAC373 are combined with 25 μ l of 10X Bam HI restriction enzyme
buffer, 20 units of Bam HI and sterile distilled water to bring the volume to 250 μ l, and
incubated at 37° for at least 3 hours. After the plasmid is digested. It is dephosphorylated
by adding one unit of calf intestinal alkaline phosphatase (CAP)/ μ g of DNA and incubated
30 for 30 min. at 37°C . The CAP is then inactivated by adding EDTA to 25 mM and SDS to
0.5% and incubated at 65°C for 15 min. The solution is then extracted with an equal volume
of phenol/chloroform/ isoamyl alcohol (25:24:1).

The aqueous phase is collected, and 125 μ l of 7.5 M ammonium acetate and 800 μ l
of 95% ethanol are added and mixed. The DNA is precipitated at -70°C for 10 min. The
35 precipitate is then collected by centrifugation by centrifugation at 12,000 rpm for 10 min.
The pellet is rinsed with (-20°C) 90% ethanol. The ethanol is then removed and the DNA
is resuspended in 50 μ l of 10 mM Tris-HCl, 1 mM EDTA, pH 7.6 (1X TE).

1

Example 32Ligation of PEDNF to pAC373

5

A purified DNA fragment containing the PEDNF gene is ligated to the transfer vector. About 200 ng of digested pAC373 are mixed with an equal molar concentration of PEDNF containing fragment. Ligation buffer (50 mM Tris-HCl, pH 7.4, 10 mM MgCl₂, 10 mM dithiothreitol, 0.5 mM spermidine, 2 mM ATP, 2.5 mM hexamine cobalt chloride and 20 µg/ml BSA) and 10 units of T4 DNA ligase is added. Water is added to a total volume of 10 µl. The mixture is incubated at 14°C for 3 hrs.

10

Example 33Transferring Genes into the AcMNPV Genome

15

Sf9 cells are seeded into 25 cm flasks at a density of 2.0×10^6 cells/flask in Grace's Antheraea medium. The cells are allowed to attach for at least one hr. One µg of MNPV DNA is added to 2 µg of plasmid DNA, which contains the PEDNF gene. The medium is removed from the flask and replaced with 0.75 ml of Grace's medium plus 10% fetal bovine serum and antibiotics (50 µg/ml gentamycin sulfate, 2.5 µg/ml amphotericin). 7.5 ml of transfection buffer (25 mM HEPES, pH 7.1, 140 mM NaCl, 125 mM CaCl₂) is added to the DNA solution and mixed. The DNA solution is added to the Grace's medium, already in the cell culture flasks.

20

Calcium phosphate precipitates form due to the calcium chloride in the transfection buffer and the phosphate in the medium. The flasks are incubated at 27°C for 4 hrs., after which time the medium is removed from the flasks, and the cells are rinsed with fresh TNM-FH (Grace's medium plus 3.3 g/l YEASTOLATE and 3.3 g/l Lactalbumin hydrolysate, both from DIFCO LABORATORIES) plus 10% fetal bovine serum and antibiotics, as described previously, and 5 ml of TNM-FH plus 10% fetal bovine and antibiotics, as described previously, is added to the cells. The cells are incubated for 4-6 days. When the infection is at an advanced stage, the virus is plated on fresh monolayers, and the recombinant viruses are plaque-purified.

25

30

Example 34Identification of Recombinant Proteins

35

Plaque-purified virus is screened by radiolabelling. The recombinant proteins are identified on SDS-PAGE gels. Sf9 cells are seeded at 6×10^5 cells/well in a 24-well culture plate. The cells are allowed to attach for an hour. The medium is then removed and overlaid with medium containing the viral stock and incubated for 1 hr. at 27°C, after which time the viral inoculum is removed and replaced with 500 µl of complete medium. The cells are incubated for 48 hrs. at 27°C. At the end of this incubation, the medium is removed. 200

-41-

1 μ l of methionine-free Grace's medium is added to the cells, and the cells are incubated for
60 min., after which time the medium is replaced with 200 μ l of fresh methionine-free
Grace's medium to which 10 μ Ci of 35 S-methionine is added. Cells are incubated at 27°C
5 for 6 hrs. and then harvested by centrifugation. The supernatant is collected, and the cells
are resuspended in 62.5 mM Tris, pH 6.8, 2% SDS, 10% glycerol, 0.001% bromophenol
blue, and 0.1 M 2-mercaptoethanol. An equal volume of 126 mM Tris, pH 6.8, 4% SDS,
20% glycerol, 0.002% bromophenol blue, and 0.2 M 2-mercaptoethanol is added to the
supernatant. Samples are then boiled for 3 min. and then they are subjected to
10 electrophoresis and autoradiography of the gel to identify proteins secreted into the medium
and proteins that are not secreted. Controls of uninfected cells and cells infected with wild-
type virus are included.

The above descriptions of exemplary embodiments of regital pigmented epithelium
derived neurotrophic factor are for illustrative purposes. Because of variations which will be
15 apparent to those skilled in the art, the present invention is not intended to be limited to the
particular embodiments described above. The present invention may also be practiced in the
absence of any element not specifically disclosed. The scope of the invention is defined by
the following claims.

20

25

30

35

-42-

1 SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Johnson, Lincoln V.
5 Tombran-Tink, Joyce

(ii) TITLE OF INVENTION: Retinal Pigmented Epithelium
Derived Neurotrophic Factor

10 (iii) NUMBER OF SEQUENCES: 1

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Christie, Parker & Hale
(B) STREET: P.O. Box 7068
15 (C) CITY: Pasadena
(D) STATE: CA
(E) COUNTRY: USA
(F) ZIP: 91109-7068

20 (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: US
(B) FILING DATE:
(C) CLASSIFICATION:

30

(vii) PRIOR APPLICATION DATA: NONE

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Sharp Esq., Janice A.
35 (B) REGISTRATION NUMBER: 34,051
(C) REFERENCE/DOCKET NUMBER: U66:23860

-43-

- 1 (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: (818)795-5843
(B) TELEFAX: (818)577-1769
- 5 (2) INFORMATION FOR SEQ ID NO:1:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1503 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
10 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- 15 (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens
20 (D) DEVELOPMENTAL STAGE: Fetus
(F) TISSUE TYPE: Eye
- (vii) IMMEDIATE SOURCE:
(A) LIBRARY: Charon BS Library of A. Swaroop
25 (B) CLONE: PEDNF
- (ix) FEATURE:
(A) NAME/KEY: prim_transcript
(B) LOCATION: 117..1370
30
- (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 117..1370
35

-44-

1 (x) PUBLICATION INFORMATION:

(A) AUTHORS: Steele, Fintan R.

Chader, Gerald J.

Johnson, Lincoln V.

5 Tombran-Tink, Joyce

(B) TITLE: Pigmented Epithelium-Derived Factor (PEDNF):

Neurotrophic Activity and Identification as a
Unique Member of the Serine Protease Inhibitor
(SERPIN) Gene Family.

10 (C) JOURNAL: Proc. Natl. Acad. Sci. USA

(G) DATE: 1992

(K) RELEVANT RESIDUES IN SEQ ID NO:1: FROM 1 TO 1502

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

15 GGACGCTGGA TTAGAAGGCA GCAAAAAAAG ATCTGTGCTG GCTGGAGCCC CCTCAGTGTG 60
   CAGGCTTAGA GGGACTAGGC TGGGTGTGGA GCTGCAGCGT ATCCACAGGC CCCAGG      116
   ATG CAG GCC CTG GTG CTA CTC CTC TGC ATT GGA GCC CTC CTC GGG CAC      164
   Met Gln Ala Leu Val Leu Leu Leu Cys Ile Gly Ala Leu Leu Gly His
       1             5             10             15
20 AGC AGC TGC CAG AAC CCT GCC AGC CCC CCG GAG GAG GGC TCC CCA GAC      212
   Ser Ser Cys Gln Asn Pro Ala Ser Pro Pro Glu Glu Gly Ser Pro Asp
           20             25             30
   CCC GAC AGC ACA GGG GCG CTG GTG GAG GAG GAG GAT CCT TTC TTC AAA      260
   Pro Asp Ser Thr Gly Ala Leu Val Glu Glu Glu Asp Pro Phe Phe Lys
25           35             40             45
   GTC CCC GTG AAC AAG CTG GCA GCG GCT GTC TCC AAC TTC GGC TAT GAC      308
   Val Pro Val Asn Lys Leu Ala Ala Ala Val Ser Asn Phe Gly Tyr Asp
       50             55             60
   CTG TAC CGG GTG CGA TCC AGC ATG AGC CCC ACG ACC AAC GTG CTC CTG      356
30 Leu Tyr Arg Val Arg Ser Ser Met Ser Pro Thr Thr Asn Val Leu Leu
   65             70             75             80
   TCT CCT CTC AGT GTG GCC ACG GCC CTC TCG GCC CTC TCG CTG GGA GCG      404
   Ser Pro Leu Ser Val Ala Thr Ala Leu Ser Ala Leu Ser Leu Gly Ala
           85             90             95

```

35

-45-

1	GAC GAG CGA ACA GAA TCC ATC ATT CAC CGG GCT CTC TAC TAT GAC TTG	452
	Asp Glu Arg Thr Glu Ser Ile Ile His Arg Ala Leu Tyr Tyr Asp Leu	
	100 105 110	
	ATC AGC AGC CCA GAC ATC CAT GGT ACC TAT AAG GAG CTC CTT GAC ACG	500
5	Ile Ser Ser Pro Asp Ile His Gly Thr Tyr Lys Glu Leu Leu Asp Thr	
	115 120 125	
	GTC ACT GCC CCC CAG AAG AAC CTC AAG AGT GCC TCC CGG ATC GTC TTT	548
	Val Thr Ala Pro Gln Lys Asn Leu Lys Ser Ala Ser Arg Ile Val Phe	
	130 135 140	
10	GAG AAG AAG CTG CGC ATA AAA TCC AGC TTT GTG GCA CCT CTG GAA AAG	596
	Glu Lys Lys Leu Arg Ile Lys Ser Ser Phe Val Ala Pro Leu Glu Lys	
	145 150 155 160	
	TCA TAT GGG ACC AGG CCC AGA GTC CTG ACG GGC AAC CCT CGC TTG GAC	644
	Ser Tyr Gly Thr Arg Pro Arg Val Leu Thr Gly Asn Pro Arg Leu Asp	
15	165 170 175	
	CTG CAA GAG ATC AAC AAC TGG GTG CAG GCG CAG ATG AAA GGG AAG CTC	692
	Leu Gln Glu Ile Asn Asn Trp Val Gln Ala Gln Met Lys Gly Lys Leu	
	180 185 190	
	GCC AGG TCC ACA AAG GAA ATT CCC GAT GAG ATC AGC ATT CTC CTT CTC	740
20	Ala Arg Ser Thr Lys Glu Ile Pro Asp Glu Ile Ser Ile Leu Leu Leu	
	195 200 205	
	GGT GTG GCG CAC TTC AAG GGG CAG TGG GTA ACA AAG TTT GAC TCC AGA	788
	Gly Val Ala His Phe Lys Gly Gln Trp Val Thr Lys Phe Asp Ser Arg	
	210 215 220	
25	AAG ACT TCC CTC GAG GAT TTC TAC TTG GAT GAA GAG AGG ACC GTG AGG	836
	Lys Thr Ser Leu Glu Asp Phe Tyr Leu Asp Glu Glu Arg Thr Val Arg	
	225 230 235 240	
	GTC CCC ATG ATG TCG GAC CCT AAG GCT GTT TTA CGC TAT GGC TTG GAT	884
	Val Pro Met Met Ser Asp Pro Lys Ala Val Leu Arg Tyr Gly Leu Asp	
30	245 250 255	
	TCA GAT CTC AGC TGC AAG ATT GCC CAG CTG CCC TTG ACC GGA AGC ATG	932
	Ser Asp Leu Ser Cys Lys Ile Ala Gln Leu Pro Leu Thr Gly Ser Met	
	260 265 270	
	AGT ATC ATC TTC TTC CTG CCC CTG AAA GTG ACC CAG AAT TTG ACC TTG	980
35	Ser Ile Ile Phe Phe Leu Pro Leu Lys Val Thr Gln Asn Leu Thr Leu	
	275 280 285	

-46-

1	ATA GAG GAG AGC CTC ACC TCC GAG TTC ATT CAT GAC ATA GAC CGA GAA	1028
	Ile Glu Glu Ser Leu Thr Ser Glu Phe Ile His Asp Ile Asp Arg Glu	
	290 295 300	
	CTG AAG ACC GTG CAG GCG GTC CTC ACT GTC CCC AAG CTG AAG CTG AGT	1076
5	Leu Lys Thr Val Gln Ala Val Leu Thr Val Pro Lys Leu Lys Leu Ser	
	305 310 315 320	
	TAC GAA GGC GAA GTC ACC AAG TCC CTG CAG GAG ATG AAG CTG CAA TCC	1124
	Tyr Glu Gly Glu Val Thr Lys Ser Leu Gln Glu Met Lys Leu Gln Ser	
	325 330 335	
10	TTG TTT GAT TCA CCA GAC TTT AGC AAG ATC ACA GGC AAA CCC ATC AAG	1172
	Leu Phe Asp Ser Pro Asp Phe Ser Lys Ile Thr Gly Lys Pro Ile Lys	
	340 345 350	
	CTG ACT CAG GTG GAA CAC CGG GCT GGC TTT GAG TGG AAC GAG GAT GGG	1220
	Leu Thr Gln Val Glu His Arg Ala Gly Phe Glu Trp Asn Glu Asp Gly	
15	355 360 365	
	GCG GGA ACC ACC CCC AGC CCA GGG CTG CAG CCT GCC CAC CTC ACC TTC	1268
	Ala Gly Thr Thr Pro Ser Pro Gly Leu Gln Pro Ala His Leu Thr Phe	
	370 375 380	
	CCG CTG GAC TAT CAC CTT AAC CAG CCT TTC ATC TTC GTA CTG AGG GAC	1316
20	Pro Leu Asp Tyr His Leu Asn Gln Pro Phe Ile Phe Val Leu Arg Asp	
	385 390 395 400	
	ACA GAC ACA GGG GCC CTT CTC TTC ATT GGC AAG ATT CTG GAC CCC AGG	1364
	Thr Asp Thr Gly Ala Leu Leu Phe Ile Gly Lys Ile Leu Asp Pro Arg	
	405 410 415	
25	GGC CCC TAATATCCCA GTTTAATATT CCAATACCCCT AGAAGAAAAC CCGAGGGACA	1420
	Gly Pro	
	GCAGATTCCA CAGGACACGA AGGCTGCCCC TGTAAGGTTT CAATGCATAC AATAAAAGAG	1480
	CTTTATCCCT AAAAAAAAAA AAA	1503

30

35

-47-

1 WHAT IS CLAIMED IS:

1. A purified retinal pigmented epithelium derived neurotrophic factor (PEDNF) composition having the amino acid sequence identified in SEQ ID NO 1.
2. A composition as recited in claim 1 wherein the composition comprises at least
5 50% by weight, with respect to the total protein, PEDNF.
3. A composition as recited in claim 1 wherein the PEDNF is a glycoprotein with a molecular weight of 50,000 to 55,000 molecular weight and a pI of 3.9 to 7.2
4. A composition as recited in claim 1 wherein the PEDNF is capable of inducing the extension of arborizing processes and neuronal differentiation.
- 10 5. A composition as recited in claim 1 wherein the PEDNF is a serine proteinase inhibitor.
6. A composition as recited in claim 1 wherein the polypeptide induces neuronal differentiation when applied to embryonic or immortalized cells.
7. A purified serine proteinase inhibitor derived from retinal pigmented
15 epithelium.
8. A purified serine proteinase inhibitor as recited in claim 7 wherein the serine proteinase exhibits neurotrophic activity.
9. A purified serine proteinase inhibitor as recited in claim 7 wherein the amino-acid composition of the serine proteinase is:
- | | | |
|----|-----|----|
| 20 | Asp | 25 |
| | Asn | 11 |
| | Thr | 28 |
| | Ser | 38 |
| | Glu | 26 |
| | Gln | 15 |
| | Pro | 29 |
| 25 | Gly | 23 |
| | Ala | 25 |
| | Val | 25 |
| | Met | 7 |
| | Ile | 22 |
| | Leu | 57 |
| | Tyr | 10 |
| | Phe | 18 |
| 30 | His | 8 |
| | Lys | 27 |
| | Trp | 3 |
| | Arg | 18 |
10. A composition as recited in claim 7 having the amino acid sequence identified in SEQ ID NO 1.
- 35 11. A purified serine proteinase inhibitor as recited in claim 7 wherein the protein comprises a glycoprotein with a molecular weight of 50,000 to 55,000 molecular weight and a pI of 3.9 to 7.2.

- 1 12. A method for purifying a retinal pigmented epithelium neurotrophic factor (PEDNF) comprising:
- providing an impure protein fraction containing PEDNF;
- applying the impure protein fraction containing PEDNF to a cation-exchange
- 5 chromatography medium;
- washing the cation-exchange chromatography medium to elute any unbound proteins;
- eluting PEDNF from the cation-exchange chromatography medium; and
- collecting the PEDNF containing fractions to provide a cation-exchange
- 10 chromatography-purified retinal pigmented PEDNF.
13. A method as recited in claim 12 wherein the impure protein fraction containing PEDNF comprises retinal pigmented epithelium conditioned media.
14. A method as recited in claim 12 wherein the impure protein fraction containing PEDNF comprises an extract of an organism transformed with a recombinant DNA molecule
- 15 comprising the PEDNF gene in a form capable of expression of the PEDNF gene.
15. A method as recited in claim 12 wherein the impure protein fraction containing PEDNF comprises a medium in which an organism, transformed with a recombinant DNA molecule comprising the PEDNF gene in a form capable of expression of the PEDNF gene, is grown.
- 20 16. A method as recited in claim 12 further comprising ammonium sulfate precipitation of the impure protein fraction containing PEDNF.
17. A method as recited in claim 16 wherein contaminating proteins are precipitated from the impure protein fraction containing PEDNF by bringing the impure protein fraction containing PEDNF to 50% to 60% saturation with ammonium sulfate to provide a 50% to
- 25 60% ammonium sulfate protein fraction.
18. A method as recited in claim 16 wherein PEDNF is precipitated from the impure protein fraction containing PEDNF by bringing the impure protein fraction containing PEDNF to 70% to 80% saturation with ammonium sulfate to provide a 70% to 80% ammonium sulfate protein fraction.
- 30 19. A method as recited in claim 16 wherein PEDNF is precipitated from the 50% to 60% ammonium sulfate fraction by bringing the 50% to 60% ammonium sulfate protein fraction to 70% to 80% saturation with ammonium sulfate to provide a 50% to 80% ammonium sulfate protein fraction.

- 1 20. A method as recited in claim 12 further comprising:
 applying the cation-exchange chromatography purified PEDNF protein fraction
 to a size-exclusion chromatography medium;
 eluting the proteins from the size-exclusion chromatography medium; and
5 collecting the PEDNF-containing fractions.
21. A method for purifying a retinal pigmented epithelium neurotrophic factor
 (PEDNF) comprising:
 providing an impure protein fraction containing; and
 purifying the PEDNF further by a method selected from the group consisting
10 of ammonium sulfate precipitation, cation-exchange chromatography, size-exclusion
 chromatography, and combinations thereof.
22. A method as recited in claim 21 wherein the purified polypeptide comprises at
 least 50% by weight, with respect to the total protein of a single polypeptide species.
23. A method as recited in claim 21 wherein the purified polypeptide induces
15 neuronal differentiation when applied to embryonic or immortalized cells.
24. A recombinant DNA molecule comprising a gene encoding a retinal pigmented
 epithelium derived neurotrophic factor (PEDNF) having the DNA sequence in SEQ ID NO
 1.
25. A recombinant DNA molecule as recited in claim 24 further comprising a
20 promoter region to direct the transcription of the PEDNF gene to form a PEDNF mRNA.
26. A recombinant DNA molecule as recited in claim 24 further comprising a
 polyadenylation region to direct the processing of the PEDNF mRNA.
27. A recombinant DNA molecule as recited in claim 24 further comprising
 ribosomal binding sites for directing the translation of the PEDNF mRNA.
- 25 28. A recombinant DNA molecule comprising a gene encoding a PEDNF having
 the amino acid sequence in SEQ ID NO 1.
29. An organism transformed with a recombinant DNA molecule comprising a
 retinal pigmented epithelium derived neurotrophic factor (PEDNF) gene having a DNA
 sequence identified in SEQ ID NO 1.
- 30 30. An organism as recited in claim 29 wherein the recombinant DNA molecule
 further comprises a promoter region, suitable for recognition by the organism transformed
 with the recombinant DNA molecule to direct the transcription of the PEDNF gene, to form
 a PEDNF mRNA.


-50-

- 1 31. An organism as recited in claim 29 wherein the recombinant DNA molecule further comprises a polyadenylation region, suitable for recognition by the organism transformed with the recombinant DNA molecule to direct the processing of the PEDNF gene, to form a PEDNF mRNA.
- 5 32. An organism as recited in claim 29 wherein the recombinant DNA molecule further comprises a ribosomal binding site, suitable for recognition by the organism transformed with the recombinant DNA molecule, to direct the translation of the PEDNF mRNA.
- 10 33. A method of treating tumors which comprises administering retinal epithelium derived neurotrophic factor (PEDNF).
34. A method as recited in claim 33 wherein the PEDNF is administered at a concentration of 1 to 100 $\mu\text{g/ml}$ 1 to 10 times per day.
35. A method of treating ocular disease which comprises administering retinal epithelium derived neurotrophic factor (PEDNF).
- 15 36. A method as recited in claim 35 wherein the PEDNF is administered at a concentration of 1 to 100 $\mu\text{g/ml}$ 1 to 10 times per day.
37. A method of treating nerve damage which comprises administering retinal epithelium derived neurotrophic factor (PEDNF).
38. A method as recited in claim 37 wherein the PEDNF is administered at a concentration of 1 to 100 $\mu\text{g/ml}$ 1 to 10 times per day.
- 20 39. A method of treating conditions resulting from the activity of serine proteases which comprises administering retinal epithelium derived neurotrophic factor (PEDNF) wherein the conditions are selected from the group consisting of excessive of unwanted blood coagulation, thrombosis, bacterial infection, parasitic infection, elastosis, vascular disorders involving fibrinoid formation, coagulation disorders, arteriosclerosis, ischemia, arthroses
- 25 diabetes, emphysema, arthritis, septic shock, lung diseases, excessive complement activation, ulcers, ulcerative colitis, pancreatitis, psoriasis, fibrinolytic disease, arthropathy, bone resorption, hypertension, congestive heart failure, cirrhosis, or allergy caused by proteases.
40. A method as recited in claim 39 wherein the PEDNF is administered at a concentration of 1 to 100 $\mu\text{g/ml}$ 1 to 10 times per day.
- 30

35

1/3

A B C D E

*Fig. 1*

2/3




Fig. 2

3/3




Fig. 3

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/05358

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) :C07K 13/00; C07H 15/12; A61K 37/02
US CL :530/399, 413; 536/23.5; 435/252.3; 514/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/399, 413; 536/23.5; 435/252.3; 514/12

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Sequence Search, Dialog, APS
search terms:retinal epithelial, neurotrophic factor

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — Y	PROC. NATL. ACAD. SCI. USA, Volume 90, issued February 1992, F.R. Steele, "Pigment epithelium-derived factor: Neurotrophic activity and identification as a member of the serine protease inhibitor gene family, pages 1526-1530, see entire document, especially figure 2.	1, 4, 5, 7-10, 24-28 ----- 2, 3, 6, 11-23, 29-40
A	DEVELOPMENTAL BIOLOGY, Volume 148, issued 1991, C.M. Park et al, "Induction of Retinal Regeneration <u>in Vivo</u> by Growth Factors," pages 322-333.	1-4

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be part of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 September 1993

Date of mailing of the international search report

14 SEP 1993

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

SHELLY GUEST CERMAK

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-0196